

Evidence for linkage of familial Diamond-Blackfan anemia to chromosome 8p23.3-p22 and for non-19q non-8p disease

Hanna Gazda, Jeffrey M. Lipton, Thiébaud-Noël Willig, Sarah Ball, Charlotte M. Niemeyer, Gil Tcherna, Naria Mohandas, Mark J. Daly, Anna Ploszynska, Karen A. Orfali, Adrianna Vlachos, Bertil E. Glader, Roma Rokicka-Milewska, Akira Ohara, David Baker, Dagmar Pospisilova, Allison Webber, David H. Viskochil, David G. Nathan, Alan H. Beggs, and Colin A. Sieff

Diamond-Blackfan anemia (DBA) is a rare congenital hypoplastic anemia that usually presents early in infancy and is inherited in 10% to 20% of cases. Linkage analysis has shown that DBA in many of both dominant and recessive DBA families mapped to chromosome 19q13.2 leading to the cloning of a gene on chromosome 19q13.2 that encodes a ribosomal protein, RPS19. However, subsequently, mutations of the RPS19 gene have only been identified in 25% of all patients with DBA. This study analyzed 14 multiplex

DBA families, 9 of which had 19q13.2 haplotypes inconsistent with 19q linkage. A genome-wide search for linked loci suggested the presence of a second DBA locus in a 26.4-centimorgan (cM) interval on human chromosome 8p. Subsequently, 24 additional DBA families were ascertained and all 38 families were analyzed with additional polymorphic markers on chromosome 8p. In total, 18 of 38 families were consistent with linkage to chromosome 8p with a maximal LOD score with heterogeneity of 3.55 at D8S277 assum-

ing 90% penetrance. The results indicate the existence of a second DBA gene in the 26.4-cM telomeric region of human chromosome 8p23.3-p22, most likely within an 8.1-cM interval flanked by D8S518 and D8S1825. Seven families were inconsistent with linkage to 8p or 19q and did not reveal mutations in the RPS19 gene, suggesting further genetic heterogeneity. (Blood. 2001;97:2145-2150)

© 2001 by The American Society of Hematology

Introduction

Diamond-Blackfan anemia (DBA) is a rare congenital hypoplastic anemia that usually presents early in infancy. The disease is characterized by a moderate to severe normochromic, often macrocytic anemia, reticulocytopenia, and selective erythroblastopenia in the bone marrow.^{1,2} Although most cases are apparently sporadic, 10% to 20% show clear patterns of inheritance, most commonly autosomal dominant.¹ DBA is markedly heterogeneous clinically, and within affected families anemia may be mild or absent in some individuals, with only subtle indications of the erythroid abnormality such as increased mean corpuscular volume (MCV) or erythrocyte adenosine deaminase activity (eADA) or both.³ More than 40% of patients have congenital abnormalities, particularly of the upper limb and craniofacial regions.^{1,4-6} Recently, linkage analysis of 29 European families showed that DBA in 26 of both dominant and recessive families mapped to chromosome 19q.⁷ Identification of a patient with a translocation led to the cloning of a gene on chromosome 19q13.2 that encodes a ribosomal protein, RPS19.^{8,9} However, mutations of the *RPS19* gene have only been identified in

25% of cases,^{8,10} suggesting a greater degree of genetic heterogeneity than expected from the initial linkage studies.^{7,11} In this report we provide evidence for the existence of a second *DBA* gene on chromosome 8p, and also for further genetic heterogeneity.

Patients, materials, and methods

Patients

Thirty-eight families with DBA from the United States, Poland, Germany, Japan, Australia, France, the Czech Republic, and the United Kingdom formed the basis of the study. Initially we ascertained DNA samples from 14 families (P1-19; Figure 1). We analyzed them for linkage to chromosome 19q and submitted them for a genome-wide screen. Subsequently these families, together with 24 additional families ascertained later (P20-49; Figure 1), were tested for linkage to candidate regions on chromosome 8. Informed consent was obtained under protocols in effect at

From the Division of Pediatric Hematology and Oncology, Dana Farber Cancer Institute and Children's Hospital, Division of Genetics, Children's Hospital and Department of Pediatrics, Harvard Medical School, Boston, Massachusetts; Division of Pediatric Hematology/Oncology, Mount Sinai School of Medicine, New York, New York; Department of Hematology, Hôpital Bicêtre, Assistance Publique-Hôpitaux de Paris et Faculté de Médecine, Paris, France; Lawrence Berkeley National Laboratory, Berkeley, California; Department of Haematology, St. George's Hospital Medical School, London, England; Children's Hospital, University of Freiburg, Germany; Whitehead Institute for Biomedical Research, Massachusetts Institute of Technology, Cambridge, Massachusetts; Department of Pediatric Hematology and Oncology, University of Gdansk, Poland; Division of Pediatric Hematology/Oncology, Stanford University School of Medicine, Stanford, California; Department of Pediatric Hematology/Oncology, Warsaw School of Medicine, Warsaw, Poland; Department of Pediatrics, Toho University School of Medicine, Tokyo, Japan; Department of Pediatric Haematology/Oncology, Princess Margaret Hospital for Children, Perth, Australia; Department of Pediatrics, University Hospital,

Olomouc, Czech Republic; Division of Medical Genetics, University of Utah School of Medicine, Salt Lake City, UT.


Submitted June 19, 2000; accepted November 22, 2000.

Supported by grants from the Daniella Maria Arturi Foundation, the Diamond Blackfan Anemia Foundation, the Association Française contre les Myopathies, Généthron, National Institutes of Health (NIH) AR44345, DK 26263, Direction de la Recherche Clinique (CRC 95183) and the Max Reinhardt Charitable Trust. H.G. was supported by F05 TW05395 from the Fogarty International Center, NIH, and A.H.B. by NIH KO2 award AR02026.

Reprints: Hanna Gazda, Pediatric Oncology, Rm M615, Dana Farber Cancer Institute, 44 Binney St, Boston, MA 02115.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 U.S.C. section 1734.

© 2001 by The American Society of Hematology



Children's Hospital, Boston, Mount Sinai Medical Center, New York, or each country of origin.

The affection status of each family member was determined by the primary hematologist in the country of origin and confirmed by our review of the clinical and laboratory data. Diagnosis was based on accepted criteria for DBA, namely, normochromic anemia in infancy or early childhood, reticulocytopenia, normal leukocyte and platelet counts, and decreased to absent bone marrow erythroid precursors. One family (P18; Figure 1) had 2 affected members with typical DBA phenotype, but normal bone marrow erythropoiesis, as described in 5% of cases.¹ All members of the DBA families were examined for congenital malformations and macrocytosis; eADA and hemoglobin F (HbF) were determined in most but not all families.

Diamond-Blackfan anemia is markedly heterogeneous with respect to the presence or absence of congenital abnormalities, the response to treatment, and evolution of the disease over time. Moreover, within families containing individuals with typical DBA, silent or mild expressions of the disease such as mild anemia or increased MCV or eADA or both have been reported in apparently unaffected family members³; indeed, in some families the *RPS19* mutation cosegregates with affected individuals and with other family members who are macrocytic or in whom eADA is elevated more than 3 SDs above the mean value.¹⁰ On clinical grounds we distinguished 2 different types of kindreds among all the multiplex DBA families. The majority of families comprised 2 or more affected individuals who presented with classical DBA with anemia, reticulocytopenia, and erythroblastopenia in the bone marrow. Coexistence of mild and severe forms of anemia in 3 families (P16, P23, and P34; Figure 1) was observed. The individuals who presented with mild anemia in these pedigrees were coded as affected in the linkage study, because they were offspring/siblings and parents of severely affected patients. Moreover, in one pedigree (P23; Figure 1), 5 affected individuals, including the member with mild anemia, presented with increased MCV (macrocytosis) and elevated eADA. This family has been described elsewhere.¹² A second group of 12 pedigrees (P14, 15, 17, 18, 22, 30, 37, 39, 43, 45, 48, 49; Figure 1) comprised classically affected probands and one or 2 individuals who presented with normal or slightly decreased hemoglobin level but with unexplained macrocytosis or highly elevated eADA or both, more than 3 SD above the mean value. These individuals were designated as genetically affected with variable expressivity of the phenotype.³ A number of affected individuals in both groups presented with various congenital malformations such as webbed neck, thumb abnormalities, congenital heart defect, urogenital malformations, and short stature. In one patient mild mental retardation was present. Family members were designated as unaffected if they had normal red blood cell counts, no congenital malformations, and did not have an unexplained elevated eADA or MCV. Thirty-two families manifested autosomal dominant inheritance; 6 pedigrees (P21, 32, 38, 44, 45, 47; Figure 1) with 2 severely affected children had parents with no clinical or laboratory signs of anemia and could represent either autosomal dominant inheritance with incomplete penetrance, gonadal mosaicism for a dominant mutation in one of the parents, or autosomal recessive inheritance. The paternity in these families was confirmed by extended haplotype analysis. The affection status of unaffected individuals in these families was coded as unknown.

DNA genotyping

DNA was isolated from whole blood by standard methods.¹³ A 10-centimorgan (cM) resolution genome-wide screen was carried out on DNA from 14 DBA families (28 potentially informative meioses) by the National Heart, Lung and Blood Institute Mammalian Genotype Service in Marshfield, Wisconsin, using their screening set 9.¹⁴ Polymerase chain reaction (PCR) and gel analysis of polymorphic markers from within chromosome

19q13.2 and chromosome 8p were performed with ³²P-labeled deoxycytosine triphosphate (dCTP), as described.¹³ Primer sequences and optimal conditions for PCR were obtained from Genome Data Base (GDB) (<http://gdbwww.gdb.org/>) and Généthon.¹⁵

RPS19 gene sequencing

The genomic DNA (gDNA) samples from affected individuals from all 38 studied families were amplified by PCR. Five PCR fragments spanning the 6 exons and 5' UTR were amplified using 150- to 200-ng template DNA in 50- μ L reactions with Taq DNA polymerase (Qiagen, Valencia, CA). PCR products were purified using a High Pure PCR Product Purification Kit (Boehringer Mannheim, Indianapolis, IN). The PCR primers and additional internal primers were used for fluorescent automated DNA sequencing, performed using Applied Biosystem 373 or 377 DNA sequencer and ABI Big Dye Terminator sequencing kits (Perkin Elmer, Foster City, CA). The DNA sequence was analyzed using Sequencher program software (GeneCodes, version 3.1).

Statistical analysis

The GENEHUNTER program (version 1.1) was used for parametric 2- and multipoint LOD score calculation¹⁶ with the assumption of autosomal dominant pattern of inheritance. Varied penetrance values were used (0.7, 0.8, 0.85, 0.9, and 0.95), with an assumed disease-gene frequency of 0.000001 (that is, disease incidence 2/1 000 000 births) and no sex difference with reference to disease frequency. Changing the disease gene frequency by several orders of magnitude had minimal influence on the results. The allele frequencies for each polymorphism, locus order, and the genetic and physical distances between them were obtained from GDB, NCBI (<http://www.ncbi.nlm.nih.gov/>) and Généthon.

Results

Chromosome 19q linkage

The PCR primers were used to amplify polymorphic markers on chromosome 19q13.2 (D19S197, LIPE, D19S408, D19S178) known to be linked to the DBA phenotype.⁷ *RPS19* is located between D19S197 and LIPE.⁸ Haplotype data for 5 of 14 initially studied families (Table 1) were consistent with linkage to chromosome 19q13.2 (P5, 7, 8, 10, 14; Figure 1). Haplotype analysis performed in the other 9 families (P1, 2, 4, 11, 15, 16, 17, 18, 19; Figure 1, Table 1) were not consistent with linkage to this region.

Genomewide screen and linkage to chromosome 8p

Because these data indicated that the locus on chromosome 19q did not explain many of our cases, a genome-wide search was performed for linked loci in all 14 pedigrees. Haplotype analysis revealed a subgroup of 8 pedigrees consistent with linkage to chromosome 8p (P1, 2, 4, 5, 11, 14, 17, 18; Figure 1); 6 pedigrees were not consistent with linkage to this region (P7, 8, 10, 15, 16, 19; Figure 1). Two of 8 families consistent with linkage to chromosome 8p were also consistent with linkage to chromosome 19q (P5, 14; Figure 1). The genome-wide search showed the highest number of pedigrees consistent with linkage to chromosome 8p, suggesting the possibility of a second *DBA* gene in that

Figure 1. Chromosomes 19q and 8p haplotypes. Structure and affected status of 38 multiplex DBA pedigrees (P). Solid symbols represent affected individuals, question mark symbols (?) represent possibly affected family members, clear symbols represent unaffected individuals. Pedigree numbers are coded to indicate that data are consistent with linkage to 8p (rectangle), 19q (oval), 8p and 19q (rectangle + oval), or inconsistent with linkage to both 8p and 19q (bold underlined) or inconclusive (parentheses), assuming 100% penetrance. The most likely haplotypes for markers D19S197, D19S408, LIPE, D19S178 and D8S518, D19S277, D8S1819, D8S439, D8S1825, and D8S1130 were created by the Cyrillic program (version 2.1).

Table 1. Results of 19q and 8p haplotype analysis and *RPS19* gene sequencing in the 38 DBA pedigrees

	UPN (P)	19q haplotype	8p haplotype	<i>RPS19</i> gene mutation	DBA phenotype in the pedigree			
					A	MCV	eADA	CA
A	1	-	+	-	SA	H	↑	Webbed neck, duplicated thumb, congenital heart defect
	2	-	+	-	SA	N	ND	Double ureter
	4	-	+	-	SA	N	ND	None
	5	+	+	-	SA	N	ND	Short stature
	7	+	-	-	SA	N	ND	Webbed neck, mental retardation
	8	+	-	-	SA	N	ND	None
	10	+	-	-	SA	H	ND	High arched palate
	11	-	+	-	SA	H	↑	Abnormal toes
	14	+	+	exon 5, T392G	SA	N	↑	None
	15	-	-	-	SA	H	↑	Extra lobe of kidney
	16	-	-	-	SA, MA	H	↑	Abnormal thumb
	17	-	+	-	SA	H	N	None
	18	-	+	-	SA	H	ND	None
	19	-	-	-	SA	N	ND	None
B	20	+	-	-	SA	H	ND	None
	21	No vertical transmission		-	SA	N	↑	None
	22	-	-	-	SA	H	↑	None
	23	+	-	exon 4, G185A	SA, MA	H	↑	Short stature
	25	+	-	-	SA	H	ND	Minor myelomeningocele
	26	-	+	-	SA	N	↑	None
	27	-	-	-	SA	N	ND	None
	28	+	+	-	SA	H	↑	None
	30	-	+	-	SA	N	↑	None
	32	No vertical transmission		-	MA	N	ND	Short stature, mental retardation
	34	-	+	-	SA, MA	H	ND	Hypertelorism, blue sclera
	37	-	+	-	SA	H	↑	Double kidney
	38	No vertical transmission		exon 3, G167A	SA	N	↑	None
	39	-	+	-	SA	N	↑	None
	40	+	+	-	SA	H	↑	None
	41	Inconclusive		-	SA	H	↑	Tripharyngeal thumbs
	42	-	-	-	SA	H	↑	DBA facies
	43	-	+	-	SA	H	↑	Growth retardation, abnormal toes, bifid thumbs, pes cavus, facies
	44	No vertical transmission		-	SA	H	↑	DBA facies, growth retardation
	45	Inconclusive-unaffected grandparents		-	MA	H	↑	None
	46	-	+	-	SA	H	↑	None
	47	No vertical transmission		-	SA	H	↑	None
	48	-	+	-	SA	N	↑	High arched palate, DBA facies, absent radial pulse
	49	-	-	-	SA	N	↑	None

Part A contains 14 initially ascertained pedigrees; part B comprises 24 pedigrees ascertained and studied subsequently. UPN (P) indicates unique pedigree number (P); +/-, consistent/inconsistent with linkage to haplotype; A, anemia; SA, severe anemia; MA, mild anemia; H, high MCV; N, in normal range; ↑, eADA > 3 SD; CA, congenital anomalies.

region. However, the result was inconclusive because of the relatively small size and number of families studied.

To address this issue, we ascertained 24 additional pedigrees (P20-49; Figure 1), and, together with the first subset of 14 families, thus analyzed 38 pedigrees in total. Eighteen of 38 families were consistent with linkage to 8p, 14 families were inconsistent, and 6 families did not show vertical transmission of the disease; it is thus difficult to assess linkage to this region in these 6 families (Table 1). They may represent autosomal dominant inheritance with reduced penetrance, mosaicism, germline mutation in one of the parents, or an autosomal recessive pattern of inheritance.

RPS19 gene mutations

Twelve of 38 pedigrees were consistent with linkage to chromosome 19q, 19 were inconsistent with linkage to this region, and

results of linkage analysis of 7 pedigrees were inconclusive (Figure 1, Table 1). Subsequent sequence analysis of the *RPS19* gene in affected individuals from all 38 families revealed different missense mutations in 3 families. One of them is a novel mutation located in exon 5 at codon 131 (T392G; Leu → Arg) in pedigree P14. Two other mutations, detected in exon 3 at codon 56 (G167A; Arg → Gln) in P38, and in exon 4 at codon 62 (G185A; Arg → Gln) in affected individuals from pedigree P23 (Figure 1, Table 1), were previously described.^{10,17} No other mutations were found in the 6 exons or their flanking regions in the other 35 pedigrees.

Multipoint linkage analysis with heterogeneity

Because the high degree of genetic heterogeneity and small family size precluded unambiguous identification of the subset of families with the second DBA locus we used the GENEHUNTER program to analyze all studied families except the 3 with mutations in exons

Table 2. GENEHUNTER analysis of 35 DBA pedigrees

Locus (from pter)	Genetic distance (cM)	Physical distance (Mb)	Chromosome 8p; HLOD scores of 35 DBA pedigrees at $\theta = 0.0$				
			Penetrance				
			70%	80%	85%	90%	95%
			HLOD (α)	HLOD (α)	HLOD (α)	HLOD (α)	HLOD (α)
D8S264	0.7	4.42	2.00 (0.47)	2.12 (0.46)	2.20 (0.44)	2.24 (0.43)	2.29 (0.41)
D8S518	6.5	7.66	2.34 (0.48)	2.44 (0.46)	2.48 (0.45)	2.53 (0.43)	2.57 (0.41)
D8S277	8.4	8.37	3.33 (0.56)	3.45 (0.53)	3.50 (0.52)	3.55 (0.50)	3.59 (0.48)
D8S1819	10	8.77	3.29 (0.55)	3.41 (0.53)	3.47 (0.51)	3.52 (0.50)	3.57 (0.47)
D8S439		8.77	3.23 (0.55)	3.34 (0.52)	3.40 (0.51)	3.45 (0.49)	3.50 (0.47)
D8S1825	14.6	9.55	2.81 (0.52)	2.93 (0.50)	2.98 (0.50)	3.04 (0.47)	3.1 (0.45)
D8S1130	14.8	11.04	2.79 (0.52)	2.91 (0.50)	2.97 (0.49)	3.03 (0.47)	3.1 (0.45)
D8S1145	26.4	17.7	0.64 (0.36)	0.68 (0.35)	0.71 (0.35)	0.73 (0.34)	0.75 (0.33)

3, 4, and 5 of the *RPS19* gene, giving a total of 35 families. The parametric multipoint LOD score with heterogeneity (HLOD score), varied between 3.33 and 3.59 at penetrance 0.7 to 0.95 for marker D8S277, with the proportion (α) of families linked to the locus equal to 0.56 to 0.48 (Table 2, Figure 2). In other words we estimate from our data that approximately 50% of families are consistent with linkage to the region of 8p with the highest HLOD score. The HLOD score decreases to 0.64 and 0.735 for locus D8S1145 at penetrances 0.7 to 0.95 (Table 2, Figure 2), suggesting strongly that the gene resides telomeric to this marker. These results provide evidence for a second *DBA* gene in the 26.4-cM interval of human chromosome 8p23.3-p22.

Evidence for non-19q non-8p DBA

In total, haplotype analysis did not reveal linkage of DBA to chromosome 8p in 14 of the 38 families (P7, 8, 10, 15, 16, 19, 20, 22, 23, 25, 27, 41, 42, 49; Figure 1, Table 1). Seven of these families also showed no evidence for linkage to chromosome 19q (P15, 16, 19, 22, 27, 42, 49); all of them were negative for *RPS19* mutations (Table 1) by sequencing. Thus, data on 7 pedigrees support

the existence of a non-19q non-8p gene (or genes). In 3 of these families haplotype segregation provided strong evidence for non-19q non-8p *DBA* gene location (P22, 27, 49) in that discordance of 19q and 8p haplotypes was found in affected family members.

Discussion

Our data suggest that a second gene for DBA resides in the telomeric region of human chromosome 8p23.3-p22. Statistical analysis revealed significant HLOD scores for the telomeric region of the chromosome 8p markers flanked by D8S1145 at penetrance 0.7 and higher. Estimation of the disease gene penetrance in DBA is difficult because of the high degree of clinical heterogeneity in this disease; however, using the single ascertainment approach¹⁸ in pedigrees that show vertical disease transmission we calculated that the disease penetrance is approximately 0.9. This result supports the significance of the HLOD score calculated at a penetrance of 0.9 (Figure 2).

Additionally we found recombinations in affected individuals in 5 of the 8p-linked families (P1, 4, 11, 17, 43) within that region. Recombinations in pedigrees P17 and P43 occurred with marker D8S518, in families P1 and P4 with D8S1145, and in pedigree P11 with marker D8S1825, suggesting that the potential critical region is flanked by the polymorphic markers D8S518 and D8S1825. These data suggested that the second *DBA* gene is possibly located within an 8.1-cM interval that contains approximately 1.89 Mb DNA and maps to chromosome 8p23.2-p23.1 (GDB html as above).

In total, haplotype analysis identified 18 families that were consistent with linkage to an 8p locus. Two of these families (P17, 18) include both severely affected members and individuals with only an increased MCV (> 3 SD above mean value) but no history of anemia. Unexplained macrocytosis in DBA kindreds in which clinically affected members also have an increased MCV suggests strongly that macrocytic individuals carry a *DBA* gene mutation and have variable expressivity of the phenotype. Variable expressivity was described in 2 families each with 2 clinically affected half-siblings and an unaffected common parent, who was therefore an obligatory gene carrier; the parent in each family had no symptoms of anemia, but did have an elevated MCV and HbF.^{19,20} Moreover, Willig et al reported a family with an *RPS19* mutation in individuals with classical DBA and in a clinically unaffected father who presented only with macrocytosis.¹⁰

In addition to the chromosome 8p- and 19q-linked families, we found 7 pedigrees inconsistent with linkage to either 8p or 19q, and negative for *RPS19* gene mutations (P15, 16, 19, 22, 27, 42, 49). In




Figure 2. Multipoint plot of HLOD scores against genetic distance (centimorgans). The analysis of 35 DBA families is shown, and the GENEHUNTER HLOD scores were calculated for the markers bounded by D8S264 (telomeric) and D8S1145 (centromeric). The data show significant linkage of a *DBA* locus to D8S277 at penetrance 0.7, 0.8, and 0.9, and to D8S1819 at penetrance 0.8 and 0.9.

4 of these, (P15, 16, 19, 42), possible explanations for this finding could be either genetic heterogeneity or lack of penetrance of the disorder in healthy siblings. However, in 3 families (P22, 27, 49) the segregation of 2 different haplotypes of chromosome 19q and 8p in affected family members provided strong evidence for a non-19q non-8p locus.

We found that the 19q families, the 8p families, and the non-8p non-19q families all contained pedigrees with the classical severe DBA phenotype as well as families with severe and mild anemia. Moreover, some individuals from each group had increased MCV or elevated eADA or both, that is, variable expressivity of the phenotype. Thus, with the proviso that insufficient families have been studied, our data do not show any correlations between clinical phenotype and genotype at this time.

Examination of candidate genes that map to the critical interval reveal no ribosomal protein genes known to map to 8p23.3-p22 region. Seventy-five of the 80 known mammalian ribosomal protein genes have been mapped,²¹ so it is possible that one of the 5 unmapped genes will localize to chromosome 8p. The region does not contain any obvious candidates, but one gene, *GATA4*, warrants discussion. *GATA4* is expressed during embryogenesis in extraembryonic visceral endoderm and appears to play an indirect role in the development of the primitive hemangioblasts that give rise to both endothelial and hematopoietic cells within blood islands of the

yolk sac²²⁻²⁴; it is also thought to be important in cardiac development. Targeted disruption of *GATA4* in mice is embryonic lethal,^{23,24} but *GATA4*^{-/-} embryonic stem cells can contribute to erythropoiesis in chimeric embryoid bodies.^{25,26} Thus a defect in *GATA4* might be expected to be extrinsic to the erythroid precursor. Although this is inconsistent with the data in DBA that suggest a defect intrinsic to erythroid cells,^{27,28} *GATA4* will be considered a candidate to be the *DBA2* gene in future studies.

In conclusion, we provide strong evidence for genetic heterogeneity of DBA. Our data indicate the existence of a second *DBA* gene on chromosome 8p23.3-p22, most likely within an 8.1-cM interval that contains approximately 1.89 Mb DNA and maps to chromosome 8p23.2-p23.1. Our results also indicate further genetic heterogeneity; that is, there is likely at least one more *DBA* gene in addition to those on chromosomes 19q and 8p. However, the high degree of genetic heterogeneity makes it difficult to generate statistical proof concerning the latter point. Confirmation of an 8p *DBA* locus will need to await the results of follow-up studies in additional patient populations as well as mutation studies of candidate genes in the region. The genetic data raise the possibility that proteins encoded by the different *DBA* genes interact in a common and novel pathway that affects erythropoiesis and other specific aspects of development.

References

- Alter BP, Young NS. The bone marrow failure syndromes. In: Nathan DG, Orkin HS, eds. Hematology of Infancy and Childhood. Philadelphia: Saunders; 1998:237-335.
- Lipton JM, Alter BP. Diamond Blackfan Anemia: Clinical Disorders and Experimental Models of Erythropoietic Failure. Boca Raton, FL: CRC Press; 1993:39-67.
- Willig T-N, Perignon JL, Gustavsson P, et al. High adenosine deaminase level among healthy probands of Diamond Blackfan anemia (DBA) cosegregates with the *DBA* gene region on chromosome 19q13. *Blood*. 1998;92:4422-4427.
- Ball SE, McGuckin CP, Jenkins G, Gordon-Smith EC. Diamond-Blackfan anaemia in the UK: analysis of 80 cases from a 20-year birth cohort. *Br J Haematol*. 1996;94:645-653.
- Janov AJ, Leong T, Nathan DG, Guinan EC. Diamond-Blackfan anemia: natural history and sequelae of treatment. *Medicine (Baltimore)*. 1996;75:77-78.
- Vlachos A, Boonyasari R, Abramson J, et al. Data from the Diamond Blackfan Anemia Registry (DBAR) suggest a reevaluation of current treatment practices. *Pediatr Res*. 1998;43:139A.
- Gustavsson P, Garelli E, Draptchinskaia N, et al. Identification of microdeletions spanning the Diamond-Blackfan anemia locus on 19q13 and evidence for genetic heterogeneity. *Am J Hum Genet*. 1998;63:1388-1395.
- Draptchinskaia N, Gustavsson P, Andersson B, et al. The gene encoding ribosomal protein S19 is mutated in Diamond-Blackfan anaemia. *Nat Genet*. 1999;21:169-175.
- Gustavsson P, Skeppner G, Johansson B, et al. Diamond-Blackfan anaemia in a girl with a de novo balanced reciprocal X;19 translocation. *J Med Genet*. 1997;34:779-782.
- Willig T-N, Draptchinskaia N, Dianzani I, et al. Mutations in ribosomal protein S19 gene and Diamond Blackfan anemia: wide variations in phenotypic expression. *Blood*. 1999;94:4294-4306.
- Gustavsson P, Willig T-N, van Haeringen A, et al. Diamond-Blackfan anemia: genetic homogeneity for a gene on chromosome 19q13 restricted to 1.8 Mb. *Nat Genet*. 1997;16:368-371.
- Viskochil DH, Carey JC, Glader BE, Rothstein G, Christensen RD. Congenital hypoplastic (Diamond-Blackfan) anemia in seven members of one kindred. *Am J Med Genet*. 1990;35:251-256.
- Engle EC, Kunkel LM, Specht LA, Beggs AH. Mapping a gene for congenital fibrosis of the extraocular muscles to the centromeric region of chromosome 12. *Nat Genet*. 1994;7:69-73.
- Yuan B, Vaske D, Weber JL, Beck J, Sheffield VC. Improved set of short-tandem-repeat polymorphisms for screening the human genome. *Am J Hum Genet*. 1997;60:459-460.
- Dib C, Faure S, Fizames C, et al. A comprehensive genetic map of the human genome based on 5,264 microsatellites. *Nature*. 1996;380:152-154.
- Kruglyak L, Daly MJ, Reeve-Daly MP, Lander ES. Parametric and nonparametric linkage analysis: a unified multipoint approach. *Am J Hum Genet*. 1996;58:1347-1363.
- Cmejla R, Blaškova J, Stopka T, et al. Ribosomal protein S19 gene mutations in patients with Diamond-Blackfan anemia and identification of ribosomal protein S19 pseudogenes. *Blood Cells Mol Dis*. 2000;26:124-132.
- Fraser FC, Nora JJ. *Genetics of Man*. Philadelphia: Lea & Febiger; 1975.
- Hunter RE, Hakami N. The occurrence of congenital hypoplastic anemia in half brothers. *J Pediatr*. 1972;81:346-348.
- Altman AC, Gross S. Severe congenital hypoplastic anemia: transmission from a healthy female to opposite sex step-siblings. *Am J Pediatr Hematol Oncol*. 1983;5:99-101.
- Kenmochi N, Kawaguchi T, Rozen S, et al. A map of 75 human ribosomal protein genes. *Genome Res*. 1998;8:509-523.
- Arceci RJ, King AA, Simon MC, Orkin SH, Wilson DB. Mouse *GATA-4*: a retinoic acid-inducible *GATA*-binding transcription factor expressed in endodermally derived tissues and heart. *Mol Cell Biol*. 1993;13:2235-2246.
- Kuo CT, Morrissy EE, Anadappa R, et al. *GATA4* transcription factor is required for ventral morphogenesis and heart tube formation. *Genes Dev*. 1997;11:1048-1060.
- Molkentin JD, Lin Q, Duncan SA, Olson EN. Requirement of the transcription factor *GATA4* for heart tube formation and ventral morphogenesis. *Genes Dev*. 1997;11:1061-1072.
- Bielinska M, Narita N, Heikinheimo M, Porter SB, Wilson DB. Erythropoiesis and vasculogenesis in embryoid bodies lacking visceral yolk sac endoderm. *Blood*. 1996;88:3720-3730.
- Soudais C, Bielinska M, Heikinheimo M, et al. Targeted mutagenesis of the transcription factor *GATA-4* gene in mouse embryonic stem cells disrupts visceral endoderm differentiation in vitro. *Development*. 1995;121:3877-3888.
- Tsai PH, Arkin S, Lipton JM. An intrinsic progenitor defect in Diamond-Blackfan anemia. *Br J Haematol*. 1989;73:112-120.
- Perdahi EB, Naprstek BL, Wallace WC, Lipton JM. Erythroid failure in Diamond-Blackfan anemia is characterized by apoptosis. *Blood*. 1994;83:645-650.