


# Developmental Cell

## Progressive Differentiation and Instructive Capacities of Amniotic Fluid and Cerebrospinal Fluid Proteomes following Neural Tube Closure

### Graphical Abstract



### Authors

Kevin F. Chau, Mark W. Springel,  
Kevin G. Broadbelt, ..., Hanno Steen,  
Anthony S. LaMantia, Maria K. Lehtinen

### Correspondence

[maria.lehtinen@childrens.harvard.edu](mailto:maria.lehtinen@childrens.harvard.edu)

### In Brief

Chau et al. provide an atlas of the developing ventricular system in mouse. Proteomics approaches reveal how amniotic and cerebrospinal fluid proteomes diverge into distinct fluid compartments from a common origin and that age-matched fluids support Sox2-positive identity in adjacent epithelia. CSF signaling via LIFR contributes to neuroepithelial cell self-renewal.

### Highlights

- Presents an atlas of dynamic, early ventricular system development
- AF and CSF proteomes diverge into distinct fluids from a common origin
- Age-matched AF and CSF promote Sox2-positive identity in developing epithelia
- CSF-based signaling via LIFR contributes to neuroepithelial cell self-renewal

# Progressive Differentiation and Instructive Capacities of Amniotic Fluid and Cerebrospinal Fluid Proteomes following Neural Tube Closure

Kevin F. Chau,<sup>1,2</sup> Mark W. Springel,<sup>1</sup> Kevin G. Broadbelt,<sup>1</sup> Hye-yeon Park,<sup>1,3</sup> Salih Topal,<sup>1</sup> Melody P. Lun,<sup>1,4</sup> Hillary Mullan,<sup>1</sup> Thomas Maynard,<sup>5</sup> Hanno Steen,<sup>1</sup> Anthony S. LaMantia,<sup>5</sup> and Maria K. Lehtinen<sup>1,2,3,\*</sup>

<sup>1</sup>Department of Pathology, Boston Children's Hospital, Boston, MA 02115, USA

<sup>2</sup>Program in Biological and Biomedical Sciences, Harvard Medical School, Boston, MA 02115, USA

<sup>3</sup>Program in Neuroscience, Harvard Medical School, Boston, MA 02115, USA

<sup>4</sup>Department of Pathology and Laboratory Medicine, Boston University School of Medicine, Boston, MA 02118, USA

<sup>5</sup>Department of Pharmacology and Physiology, Institute for Neuroscience, The George Washington University School of Medicine and Health Sciences, Washington, DC 20037, USA

\*Correspondence: maria.lehtinen@childrens.harvard.edu

<http://dx.doi.org/10.1016/j.devcel.2015.11.015>

## SUMMARY


After neural tube closure, amniotic fluid (AF) captured inside the neural tube forms the nascent cerebrospinal fluid (CSF). Neuroepithelial stem cells contact CSF-filled ventricles, proliferate, and differentiate to form the mammalian brain, while neurogenic placodes, which generate cranial sensory neurons, remain in contact with the AF. Using *in vivo* ultrasound imaging, we quantified the expansion of the embryonic ventricular-CSF space from its inception. We developed tools to obtain pure AF and nascent CSF, before and after neural tube closure, and to define how the AF and CSF proteomes diverge during mouse development. Using embryonic neural explants, we demonstrate that age-matched fluids promote Sox2-positive neurogenic identity in developing forebrain and olfactory epithelia. Nascent CSF also stimulates SOX2-positive self-renewal of forebrain progenitor cells, some of which is attributable to LIFR signaling. Our Resource should facilitate the investigation of fluid-tissue interactions during this highly vulnerable stage of early brain development.

## INTRODUCTION

Morphogenesis of the cerebrospinal fluid (CSF)-filled ventricular system requires a series of highly stereotyped, conserved events (Bjornsson et al., 2015; Lowery and Sive, 2009; Massarwa and Niswander, 2013; Wilde et al., 2014). Tremendous changes in ambient fluid composition and brain morphology occur at embryonic day (E) 8.5-E10.5 in the mouse embryo, yet the relationship between these changes is poorly understood. At E8.5, the neuroectoderm destined to become the mammalian brain is exposed to amniotic fluid (AF). As the anterior neural tube closes, the AF captured inside the

neural tube becomes the nascent CSF, and the entire mammalian brain subsequently develops around CSF-filled ventricular spaces (Ek et al., 2005; Zappaterra and Lehtinen, 2012). Subsequently, the neurogenic placodes that give rise to cranial sensory neurons (Moody and LaMantia, 2015) remain exposed to AF, while the developing brain is exposed to CSF. As brain development progresses, the forebrain neuroectoderm consists of neuroepithelial cells that rapidly divide to expand the neural progenitor cell pool (Bjornsson et al., 2015). Over subsequent days, the neuroepithelium becomes patterned, with well-defined gene expression domains along the anterior-posterior and dorsal-ventral axes, ultimately giving rise to the wealth of cells in distinct regions of the mammalian brain (Grove and Monuki, 2013; Rallu et al., 2002a). The progressive lineage restriction of neural progenitors and transition to radial glial progenitors of the cerebral cortex occurs during this period. While pressurization of the ventricles is essential for normal brain development to proceed (Desmond and Jacobson, 1977; Lehtinen and Walsh, 2011), little is known about the signals delivered by the AF versus nascent CSF and whether they instruct the development of adjacent epithelia during these early stages of brain development.

Cell intrinsic genetic programs regulate progenitor cell lineage restriction (Shen et al., 2006). However, extrinsic signals play important roles in specifying progenitor identity as well (Hunter and Hatten, 1995; LaMantia et al., 1993; McConnell and Kaznowski, 1991). The interplay between genetic programs (Manzini and Walsh, 2011; Shen et al., 2006), CSF (Lehtinen et al., 2011; Lun et al., 2015b), meningeal (Chatzi et al., 2013; Siegenthaler et al., 2009), and vascular (Kokovay et al., 2010; Shen et al., 2004; Tavazoie et al., 2008) signals helps coordinate and synchronize brain development at later ages. At these later stages, secreted signals in the embryonic CSF (Lehtinen et al., 2011), many of which originate from the choroid plexus (Huang et al., 2009c; Lun et al., 2015a; Yamamoto et al., 1996), instruct the proliferation and survival of polarized radial glial progenitor cells as well as adult neural stem cells, in part via signal transduction at the cilium (Higginbotham et al., 2013; Tong et al., 2014; Yeh et al., 2013). Proteomic approaches have begun to characterize



**Figure 1. Expansion of the Mouse Ventricular System following Neural Tube Closure**

(A) SOX2-positive neuroepithelial cells (green) contact the AF before neural tube closure at E8.5 (left) and the nascent CSF following neural tube closure at E9.5 (middle) and E10.5 (right). The nuclei were counterstained with Hoechst. The scale bars represent 50  $\mu$ m (left) and 200  $\mu$ m (middle and right).

(B) Representative ultrasound images of developing mouse (E8.5–E12.5). The dashed white line (left) outlines the embryo. The scale bar represents 2 mm.

(C) 3D reconstruction of ventricular system from ultrasound image sequences captured in (B) from E9.5–E12.5 with the following color representation: turquoise: lateral ventricles; yellow: third ventricle and aqueduct; and dark blue: fourth ventricle.

(D) Quantification of ventricle size based on 3D reconstructions of ultrasound image sequences (B and C, data not shown; ventricular volume ( $\mu$ l)) represented as mean  $\pm$  SEM; lateral ventricles: E9.5 =  $0.09 \pm 0.02$ ; E10.5 =  $0.77 \pm 0.12$ ; E11.5 =  $0.83 \pm 0.07$ ; E12.5 =  $1.39 \pm 0.19$ ; E13.5 =  $1.08 \pm 0.05$ ; third ventricle and aqueduct: E9.5 =  $0.31 \pm 0.03$ ; E10.5 =  $1.48 \pm 0.37$ ; E11.5 =  $1.33 \pm 0.15$ ; E12.5 =  $1.40 \pm 0.11$ ; E13.5 =  $1.16 \pm 0.03$ ; fourth ventricle: E9.5 =  $0.54 \pm 0.04$ ; E10.5 =  $1.63 \pm 0.22$ ; E11.5 =  $1.36 \pm 0.06$ ; E12.5 =  $1.27 \pm 0.27$ ; E13.5 =  $0.50 \pm 0.14$ ; total ventricular volume: E9.5 =  $0.94 \pm 0.04$ ; E10.5 =  $3.92 \pm 0.70$ ; E11.5 =  $3.51 \pm 0.16$ ; E12.5 =  $4.06 \pm 0.06$ ; E13.5 =  $2.74 \pm 0.09$ ; n = 3; ANOVA statistical analyses of total ventricular volume represented as (\*) p < 0.05 and (\*\*) p < 0.0001; and n = 3.

the composition of the CSF during later stages of development in different species (see Chang et al., 2015; Dziegielewska et al., 1981; Lehtinen et al., 2011; Lun et al., 2015a; Parada et al., 2005; Zappaterra et al., 2007), as well as in adult (see Dislich et al., 2015). However, around the time of neural tube closure (E8.5–E10.5), the choroid plexus does not exist (Lun et al., 2015b) and the meninges as well as the brain's vasculature are only beginning to form (Björnsson et al., 2015). The only source of soluble extrinsic cues for neural stem cells at this time is the AF and nascent CSF, which contact the apical surface of all neural stem cells. Nevertheless, it is not known whether distinct AF and nascent CSF instruct early neuroepithelial cells and very few clues exist regarding the identity of such instructive signals. Indeed, even the basic proteomic composition of early embryonic AF and nascent CSF remains undefined.

To provide a resource to formulate and answer critical questions of how early neural stem cells in the central as well as peripheral nervous system are regulated by the adjacent fluid proteomes, we developed techniques to collect AF and nascent CSF from developing embryonic mouse before and after neural tube closure. We demonstrate the dynamic capacity of these fluids to signal via cardinal morphogenetic signaling pathways including Sonic hedgehog (SHH), Bone morphogenetic proteins (BMPs), and retinoic acid (RA). Using quantitative mass spectrometry (MS) of these fluids, we define their progressive proteomic differentiation from their common AF origin at E8.5, through

E10.5 and E14.5 when distinct AF and CSF compartments exist. We developed a neural explant culture system to establish that age-matched fluid-tissue interactions promote Sox2-positive neurogenic identity in developing forebrain neuroepithelium as well as peripheral olfactory epithelium (OE). Nascent CSF promotes SOX2-positive self-renewal of forebrain neural stem cells, and we demonstrate a role for signaling via the leukemia inhibitory factor (LIF) receptor (LIFR) in this process. Thus, our comprehensive resource defines divergence between the AF and CSF proteomes and validates the functional significance of these differences using *in vitro* and *ex vivo* approaches, facilitating the investigation of fluid-tissue interactions during the initial formation of the nervous system.

## RESULTS

### Rapid Morphological Changes in Brain-Ventricular System during Early Development

In the vertebrate embryo, the transition from an open anterior neural plate contacted by AF to a closed forebrain adjacent to CSF-filled ventricles is a key morphogenetic event in brain development. At E8.5 in mouse, the neuroectoderm, which is home to SOX2-positive neuroepithelial cells, directly contacts the AF (Figure 1A). As the anterior neural tube closes, the AF captured inside the neural tube becomes the nascent CSF (Ek et al., 2005; Zappaterra and Lehtinen, 2012). By E9.5, SOX2-positive

neuroepithelial progenitor cells in the developing forebrain directly contact the CSF (Figure 1A), and the forebrain continues to expand around this CSF-filled ventricle (Figure 1A). To evaluate the influence of these morphogenetic changes for defining the CSF as a distinct entity, we performed *in utero* ultrasound imaging of live embryos (Figure 1B) and quantified regional differences in the establishment and expansion of the ventricular system during this early phase of brain development (Figures 1C and 1D). Following neural tube closure at E9, the lateral ventricles continuously expanded until E12.5 (mean ventricular volume ( $\mu\text{l}$ )  $\pm$  SEM: E9.5 =  $0.092 \pm 0.015$ ; E12.5 =  $1.392 \pm 0.193$ ;  $t$  test,  $p < 0.005$ , and  $n = 3$ ; Figures 1C and 1D), while third and fourth ventricle expansion was most prominent between E9.5–E10.5 (Figures 1C and 1D). Ventricular volume in live embryos was quite distinct from that measured in fixed tissue in a separate set of samples. Paraformaldehyde fixation caused shrinkage of the ventricles by approximately 60% (data not shown). Taken together, following its definition with the closure of the anterior neural tube, the mouse ventricular system undergoes dynamic morphological changes in a ventricle-specific manner.

#### Age-Matched AF and CSF Support Neurogenic Cell Fate

Given the striking change in fluid environment that neural tube closure imposes on the developing neuroectoderm, we tested if E8.5 AF and E10.5 CSF have the capacity to differentially instruct neuroepithelial cell fate before and after neural tube closure. We first developed methods to collect pure samples of AF and CSF at these early ages. To collect E8.5 AF, the decidua and parietal yolk sac were removed to expose the visceral yolk sac, and AF was withdrawn from the amniotic cavity (Figure 2A). The E8.5 embryo did not appear to incur damage due to AF collection (Figures S1A and S1B). AF was similarly collected from the amniotic cavity at E10.5 (Figure 2B). To collect E10.5 CSF, the extra-embryonic membranes were removed and CSF was withdrawn from the telencephalic vesicles, aqueduct, and fourth ventricle (Figure 2C). Following fluid collection, samples were verified for purity (based upon absence of red blood cells or other particulate matter) and processed (Zappaterra et al., 2013). Standard litters (10–12 embryos) yielded the following volumes: E8.5 AF = 10–15  $\mu\text{l}$ ; E10.5 AF = 180–225  $\mu\text{l}$ ; and E10.5 CSF = 30–40  $\mu\text{l}$ . CSF yield was comparable to the ventricular size calculated by ultrasound (e.g., E10.5 ventricular volume =  $3.9 \pm 0.7 \mu\text{l}/\text{embryo}$ ; Figure 1D).

We then developed a protocol for forebrain neuroectodermal explants at E8.5 and E10.5 and tested for the instructive capacity of AF and nascent CSF. Once the neuroepithelium was separated from mesenchyme and surface ectoderm (both are sources of local signals that might confound assessment of AF or CSF signaling capacity; see LaMantia et al., 2000; Tucker et al., 2008), explants were exposed to AF or CSF from different ages (Figure 2D). Heterochronic forebrain neuroectodermal explants using Sox2:EGFP knockin mice (Ellis et al., 2004) suggested that progenitor cell identity is differentially regulated by secreted signals received from E8.5 AF versus E10.5 CSF. Survival as well as the neurogenic phenotype of immature neuroectodermal cells at E8.5, as assessed by Sox2:EGFP in explants, appeared optimally supported by age-matched E8.5 AF (Figures 2E and 2F). Conversely, E10.5 CSF optimally supported E10.5 neuroectodermal cells (Figures 2G and 2H). These data demon-

strate that compartment/fluid/tissue age-matching—either via changes in osmolality or through distinct signaling capacities—contributes to the maintenance of neurogenic progenitor cell fate.

#### Dynamism of AF and CSF and Age-Dependent Instructive Capacities of AF


There is very little insight into the quantitative changes in soluble factors that occur during the time that the CSF becomes a distinct fluid compartment from the AF. Measurement of the total protein concentration in AF and nascent CSF suggested that the proteomes of these fluids are tightly regulated. The total AF and CSF protein concentrations started to decrease from E8.5–E10.5 (Figure 3A), and the total AF protein concentration continued to decrease to E14.5 (Figure 3A). In contrast, the total CSF protein concentration sharply increased from E10.5–E14.5 (Figure 3A), consistent with previous studies (Ek et al., 2005; Lun et al., 2015a). Unexpectedly the E14.5 CSF proteome appeared less complex in comparison with E8.5 AF and E10.5 CSF when analyzed by silver staining (Figure 3B). The increase in E14.5 CSF protein concentration was apparently due to robust increases in certain proteins, such as albumin, which appeared saturated by silver staining (Figure 3B) and which we verified by quantitative MS to increase approximately 2-fold between E10.5 CSF to E14.5 CSF (albumin mean spectral counts  $\pm$  SEM: E10.5 CSF =  $97.1 \pm 16.6$ ; E14.5 CSF =  $201.6 \pm 23.6$ ; Mann-Whitney,  $p < 0.05$ , and  $n = 3$ ; see also Table S1).

As the ventricular system expands during the course of early development, the CSF draws fluid from the AF, a process suggested to be driven in part by an osmolarity gradient (Alonso et al., 1998; Gato and Desmond, 2009). Despite the changes observed in total protein concentration between the fluids (Figure 3A), we did not detect differences in global osmolality between AF and CSF during this time frame (Figure 3C). These data agree with the model in which high colloid osmotic pressure, which typically constitutes a small fraction of the total osmotic pressure and is generated locally by individual or groups of proteins, drives ventricular expansion (Ek et al., 2005).

We next tested if the AF, like the CSF, harbors instructive signaling activities for adjacent tissues. We asked whether a cranial neurogenic placodal epithelium, where neural stem cells for peripheral neurons are established and maintained (Moody and LaMantia, 2015), depends selectively on the signaling capacity of the AF, which contacts these structures. We developed an E10.5 olfactory placode explant culture system (Figure 3D), which gives rise to the olfactory sensory neurons and which under normal conditions directly contacts the E10.5 AF. The neurogenic phenotype of developing E10.5 olfactory epithelial cells was optimally supported by E10.5 AF versus E8.5 AF, as assessed by Sox2:EGFP (Figures 3E and 3F), and the lateral to medial patterning of Sox2 (Rawson et al., 2010; Tucker et al., 2010) was also optimally maintained in the presence of E10.5 AF. These data demonstrate that the AF can regulate cranial placode neuroepithelial cells in an age- and tissue-specific manner.

#### Quantitative Proteomic Analyses Reveal Dynamic Transition from AF to CSF

To characterize the differentiating CSF proteome as it emerges from E8.5 AF, we performed quantitative MS using label-free



**Figure 2. Heterochronic Explants Show that Tissue-Fluid Age Matching Promotes *Sox2*-Positive Progenitors**

(A–D) Schematics of (A) E8.5 AF collection; (B) E10.5 AF collection; (C) E10.5 CSF collection; and (D) heterochronic forebrain neuroectodermal explant dissections: isolated neuroepithelium (dashed line) placed on membrane with ventricular surface down contacting AF or CSF.

(E) E8.5 explants grown on E8.5 AF or E10.5 CSF. The *Sox2:EGFP* suggests age-matched fluid best supports neural stem cell identity (E8.5 explant with E8.5 AF [n = 8] or with E10.5 CSF [n = 8]).


(F) Quantitative (q)PCR of explants cultured as in (E) suggests that E8.5 AF more favorably supports E8.5 explants compared to E10.5 explants (mean *Sox2* fold expression ± SEM: E8.5 AF = 1.63 ± 0.49; E10.5 CSF = 1.00; n = 3 of 4 pooled explants each, p = not significant, n.s., and Mann-Whitney). There were two experiments (n = 2) of four pooled explants each that were excluded from statistical analyses. While E8.5 AF supported these E8.5 explants, the experiments had undetectable *Sox2* expression in the E10.5 CSF condition, suggesting a role for age-matched AF and CSF in cell survival and demonstrating experimental variability at this age (see also Discussion).

(G) E10.5 explants grown on E8.5 AF [n = 9] or with E10.5 CSF [n = 11]. Note increased GFP expression for age-matched CSF versus non-age-matched CSF experiments.

(H) qPCR of explants cultured as in (G) show that E10.5 CSF more favorably promotes *Sox2* expression in E10.5 explants compared to E8.5 AF (mean *Sox2* fold expression ± SEM: E8.5 AF = 1.00; E10.5 CSF = 1.80 ± 0.29; n = 3 of 4 pooled explants each, p < 0.05, and Mann-Whitney). See also Figure S1.

spectral counts on E8.5 AF, E10.5 CSF, and E14.5 CSF. Consistent with the silver staining results (Figure 3B), E8.5 AF had the most complex proteome (764 proteins identified), followed by E10.5 CSF (504 proteins), while E14.5 CSF had the least complex proteome (410 proteins) (Figures 4A and 4B; Table S1). All three fluid compartments shared 225 proteins (e.g., APOA4 and FSTL1) (Figures 4A and 4B; Table S1). Beyond these shared constituents, we found 383 proteins exclusive to E8.5 AF (e.g., EEF1G and RPL6), whereas 126 and 32 were exclusive to E10.5 CSF (e.g., ACTG1 and EFNB2) and E14.5 CSF (e.g., BLBP/FABP7 and CDH13), respectively (Figures 4A and 4B; Table S1).

Unsupervised clustering using GProX (Rigbolt et al., 2011) partitioned the proteomes into six individual clusters with distinct temporal expression patterns as the fluids diverged from E8.5 AF (Figure 4C; Table S1). Cluster 1 consisted of 452 proteins that were more abundant at E8.5 AF than at later stages. Enrichment analysis using DAVID Web tool (Huang et al., 2009a, 2009b) showed that ribosomal proteins and other proteins involved in translation constituted the most enriched class of proteins in cluster 1 (Figure 4D; Table S1). The early neuroepithelium releases membrane particles (Marzesco et al., 2005). Consistent with these findings, immunostaining analyses of some ribosomal protein subunits suggested to be enriched in E8.5 AF based on our MS analyses (e.g., RPS12 and RPL11) were expressed in the E8.5 neuroepithelium at higher levels than in the E10.5 neuroepithelium (Figures S1C and S1D), strongly suggesting that ribosomal protein content in E8.5 AF arises from normal membrane shedding that occurs during this age. Cluster 2 (181 members) represented proteins that were more abundant in E10.5 CSF than in E8.5 AF and E14.5 CSF.



**Figure 3. Early AF and CSF Undergo Dynamic Changes, and Age-Matched AF Supports Olfactory Placode Development**

(A) Total protein concentration of AF decreases from E8.5–E14.5; total CSF protein concentration increases from E10.5–E14.5 (mean protein concentration [mg/ml] ± SEM: E8.5 AF = 1.99 ± 0.06; E10.5 AF = 1.73 ± 0.13; E10.5 CSF = 1.70 ± 0.14; E14.5 AF = 1.51 ± 0.05; E14.5 CSF = 2.73 ± 0.26; n = 4 for all except E10.5 CSF, n = 5; ANOVA; and \* p < 0.05 and \*\*\* p < 0.0001).

(B) Silver stain of 1 µl AF or CSF suggests highest protein complexity in E8.5 AF. The asterisks denote representative proteins enriched in E8.5 AF (red) or E14.5 CSF (green).

(C) Osmolality of CSF decreases from E8.5–E10.5 and then increases. The osmolality of AF decreases from E8.5–E14.5 (mean osmolality [mOsm/Kg] ± SEM: E8.5 AF = 303.0 ± 3.0; n = 3; E10.5 AF = 296.8 ± 2.2; n = 5; E10.5 CSF = 293.4 ± 4.4; n = 5; E12.5 AF = 292.5 ± 2.5; n = 4; E12.5 CSF = 297.0 ± 5.3; n = 3; E14.5 AF = 293.7 ± 1.2; E14.5 CSF = 298.0 ± 1.5; n = 3; ANOVA, and p < 0.05). The osmolality of E14.5 CSF is higher compared to E14.5 AF (t test and \*p < 0.05).

(D) Schematic of E10.5 OE dissections: pairs of bilateral olfactory epithelial explants (denoted by dashed lines) placed on porous membrane contacting the AF.

(E) E10.5 olfactory epithelial explants grown on E8.5 or E10.5 AF. The *Sox2:EGFP* shows age-matched fluid most favorably supports neural stem cell identity (n = 4).

(F) qPCR of E10.5 OE explants cultured as in (E) show E10.5 AF more favorably promotes *Sox2* expression in E10.5 OE explants compared to E8.5 AF (mean *Sox2* fold expression ± SEM: E8.5 AF = 1.00; E10.5 AF = 1.78 ± 0.32; n = 3 of 4 pooled explants each, p < 0.05, and Mann-Whitney).


The most enriched class within cluster 2 was glycoproteins (Figure 4D; Table S1), many of which were membrane proteins (e.g., cadherins and prominin1) and extracellular matrix (ECM) proteins (e.g., collagens, laminins, and reelin), likely reflecting the active proliferation and membrane shedding of the neuroepithelium at E10.5. Supporting these data, KEGG pathway analysis of cluster 2 showed that ECM-receptor interaction ( $p$  value =  $5.89 \times 10^{-9}$ ) and focal adhesion ( $p$  value =  $6.84 \times 10^{-6}$ ) were the two most enriched pathways within this cluster (Table S1). The third most enriched pathway was axon guidance ( $p$  value =  $4.03 \times 10^{-5}$ ; Table S1), with members such as semaphorins, ephrin receptors, ephrin-B2, and DCC (netrin-1 receptor) (Figure 4E; Table S1), reflecting the active cellular movement in the embryo at this age. Whether these activities have active roles in instructing migration, as has been suggested for CSF-SLIT in adult (Sawamoto et al., 2006), remains to be determined. A number of proteins involved in canonical signaling pathways, including BMP1, MSTN, DKK3, and TGFBR3, were also found in cluster 2 (Table S1), demonstrating that these activities are already present in E10.5 CSF, before the development of the choroid plexus. The availability of these proteins was downregulated by E14.5, suggestive of roles in early brain development. Cluster 3 (87 members) represented proteins enriched in E14.5 CSF, and the most enriched class within this cluster was proteins with

immunoglobulin domains. Taken together, these data demonstrate that the AF-to-CSF transition during neural tube closure is accompanied by dynamic changes in protein availability and putative functions.

#### Canonical Signaling Activities in AF and CSF

A number of canonical signals (e.g., SHH, BMPs, and RA) act on neural stem cells in the developing brain during the time when the CSF becomes a distinct fluid compartment (Wilde et al., 2014), and we found that these activities are tightly regulated in the nascent CSF during this time frame. First, from our MS analysis, the non-canonical SHH receptors, BOC and GAS1, were detected in E10.5 CSF (Table S1). Consistent with these data, CSF-SHH, quantified by ELISA, peaked in E10.5 CSF (Figure 5A). The fourth ventricle choroid plexus secretes SHH into the CSF at later ages (Huang et al., 2010; Lun et al., 2015a); however, the choroid plexi have not yet developed at E10.5. SHH enrichment in the nascent CSF supports a model in which secretion of signals from cells adjacent to the ventricles, and/or diffusion of signals from surrounding tissues (e.g., floor plate), contribute to the CSF proteome, akin to secretion of SEMA3B into the CSF in the spinal cord (Arbeille et al., 2015).

Previous studies have demonstrated activation of BMP signaling in apical progenitor cells along the length of the dorsal




**Figure 4. MS Reveals Dynamic Changes in CSF Proteome as It Differentiates from E8.5 AF to E14.5 CSF**

(A) Heatmap of normalized spectral counts reveals differential protein availability between E8.5 AF, E10.5 CSF, and E14.5 CSF. The unsupervised hierarchical clustering grouped the three biological replicates of each fluid together. Each replicate contains 30 µg total CSF protein pooled from multiple embryos. The spectral counts were scaled such that for each protein, the sample with the highest spectral count is set as one. The gray indicates undetected protein (i.e., 0 spectral counts). Proteins with peptides detected in at least two samples were included in the heatmap.

(B) Total of 961 proteins were identified across all samples. The Venn diagram depicts the numbers of proteins exclusive to or shared between distinct fluid compartments.

(C) Unsupervised clustering using GProX partitioned proteins into six clusters, revealing different temporal expression patterns for each cluster. The number of proteins in each cluster ( $n$ ) is indicated above each graph. There were 33 proteins that showed less than 2-fold difference in availability between the fluids and were not included in clustering. Membership indicates how well the proteins fit into the general profile of cluster.

(legend continued on next page)



(D) RA activity measured in E8.5 AF, E10.5 CSF, and E14.5 CSF as luciferase signal in clonally derived RA-sensitive cell line. The responses were compared to linear responses generated by pure RA in the same cell line (data not shown). The overall RA activity increases in E14.5 CSF (mean RA activity + SEM: E8.5 AF = 12.3 ± 2.0; E10.5 CSF = 15.3 ± 3.4; E14.5 CSF = 33.4 ± 7.4; t test,  $p < 0.01$ ;  $n = 3$ ; data represent duplicate runs of biological replicates, and  $n = 3$ , at each age).

(E) RA indicator mice at E8.5 (left) with open neural tube and E10.5 (right) with closed neural tube (arrows: presumptive, developing forebrain). The RA signaling is not detected in the forebrain neuroectoderm of the open neural tube at E8.5. The signaling in the somites and spinal cord reflects local RA production by the somites (Stavridis et al., 2010; Vermot et al., 2005). At E10.5, RA signaling activity in the forebrain and head is modest, confined to the ventro-lateral forebrain, eye, and nose, all reflecting local neural crest mesenchyme sources of RA (Haskell and LaMantia, 2005; LaMantia et al., 1993).

telencephalic ventricular surface (Doan et al., 2012; Lehtinen et al., 2011), and we found BMP signaling activity to change dynamically in the nascent CSF. BMP4 availability measured by ELISA showed a trend to increase from E8.5 AF to E14.5 CSF, but was not significant (mean BMP4 [nmol/ml] ± SEM: E8.5 AF = 745.1 ± 111.7; E10.5 CSF = 941.6 ± 201.4; E14.5 CSF = 1018.3 ± 231.4;  $n = 3$ , and not significant). We observed progressively increased phospho-SMAD1/5/8-positive staining cells along the ventricular surface during neurulation (mean forebrain phospho-SMAD1/5/8 frequency/mm ± SEM: E8.5 = 28.0 ± 1.6 [ $n = 8$ ]; E9.5 = 75.3 ± 3.2 [ $n = 5$ ]; E10.5 = 83.0 ± 2.9 [ $n = 6$ ]; E8.5 versus E9.5, t test,  $p < 0.0001$ ; E9.5 versus E10.5, t test, and  $p < 0.05$  and Figure 5B), suggesting that these cells have the ability to transduce Bmp signals from the AF and CSF. In contrast, overall BMP activity progressively decreased from E8.5 AF to E14.5 CSF, as revealed by a luciferase-based assay in a BMP-sensitive cell line (Figure 5C). Finally, RA activity was similarly low between E8.5 AF and E10.5 CSF and increased by E14.5 (Figure 5D), after choroid plexus formation, consistent with a requirement during initial neural tube formation to “protect” the anterior neural plate from the posteriorizing influence of RA (Durston et al., 1989), followed by increased RA signaling in more mature forebrain neural stem cells (Figure 5E) (Haskell and

LaMantia, 2005; LaMantia et al., 1993; Lehtinen et al., 2011; Siegenthaler et al., 2009). Together, these data show that signaling activities with established roles in brain development are dynamically available in the AF and nascent CSF and parallel changes in signaling in the neural progenitors.


### Nascent CSF Supports Progenitor Self-Renewal

We next asked if these distinct, dynamic AF and CSF proteomes could be mined as a resource for signaling activities that might influence neural stem and progenitor progression. Proteomic analyses revealed the dynamic availability of many receptors from E8.5 AF–E14.5 CSF (e.g., TGFβ, EGFR, and LIFR, representative of 41 proteins annotated as receptors by DAVID; Figures S2A–S2C; Table S1). For example, availability of the LIFR declined during the course of AF to CSF fluid maturation (LIFR mean spectral counts ± SEM: E8.5 AF = 14.8 ± 1.2; E10.5 CSF = 7.8 ± 2.8; E14.5 CSF = not detected, and  $n = 3$ ; Figure S2C; Table S1). While LIFR was not detected by MS in E14.5 CSF, immunoblotting revealed bands consistent with some LIFR availability in E14.5 CSF (Figure S2D). Soluble LIFR binds LIF and can inhibit its biological activity (Layton et al., 1992; Tomida, 2000). These observations suggest that neural tube closure could decrease access of an inhibitory factor into the CSF.

(D) Functional annotation clustering of proteins in GProX clusters 1, 2, and 3 using DAVID. The top five enriched functional clusters are shown.

(E) Proteins in GProX cluster 2 that are members of the axon guidance pathway were subjected to network analysis using GeneMania (Warde-Farley et al., 2010). The green nodes represent protein members of cluster 2, whereas the gray nodes represent related proteins. The lines connecting the nodes denote the relationship between the proteins.

See also Table S1 and Figure S1.



**Figure 6. Nascent CSF Promotes Signaling and Self-Renewal in Progenitor Cells**

(A) E10.5 forebrain shows P-STAT3 (red) activity in P-Vimentin-positive (green) progenitor cells along ventricular surface. The nuclei were counterstained with Hoechst.

(B) Higher magnification images of E10.5 forebrain show LIFR (red) expression in P-Vimentin-positive (green) progenitor cells along ventricular surface. The nuclei were counterstained with Hoechst. The scale bar represents 20  $\mu$ m.

(C) Removal of N-linked glycans from E10.5 CSF with PNGase F reveals robust bands immunoreactive for LIF.

(D) Intraventricular LIF injection (15 min; 200 ng/ml) at E10.5 stimulates P-STAT3 (red) signaling in P-Vimentin-positive (green) progenitor cells along ventricular surface. The nuclei were counterstained with Hoechst.

(E) Schematic depicting potential combinations of daughter cells in pair cell assay (progenitor: P; neuron: N; and transient/intermediate cell type co-expressing SOX2 and TUJ1: Tr).

(F) LIF stimulates SOX2-positive self-renewal (P-P) of progenitor cells at expense of differentiating cells (P-N; Tr-Tr; identities of pairs of cells represented as mean  $\pm$  SEM; P-P division: control =  $23.3 \pm 0.8$ ; LIF =  $35.9 \pm 1.6$ ; P-N division: control =  $18.8 \pm 0.9$ ; LIF =  $11.6 \pm 0.9$ ; Tr-Tr division: control =  $38.0 \pm 1.9$ ; LIF =  $28.8 \pm 2.0$ ; N-N division: control =  $20.1 \pm 2.5$ ; LIF =  $23.7 \pm 0.6$ ; n = 3; p < 0.0001; and ANOVA).

(G) E10.5 CSF (20%) stimulates SOX2-positive self-renewal (P-P) of progenitor cells at expense of differentiating cells (data presented as above in F) (P-P division: control =  $33.3 \pm 2.5$ ; CSF =  $40.2 \pm 2.2$ ; P-N division: control =  $18.2 \pm 0.9$ ; CSF =  $23.4 \pm 3.4$ ; Tr-Tr division: control =  $36.8 \pm 2.1$ ; CSF =  $26.0 \pm 0.4$ ; N-N division: control =  $12.8 \pm 0.8$ ; CSF =  $12.2 \pm 1.3$ ; n = 3; p < 0.05; and ANOVA).

(H) Interference with LIF signaling in E10.5 CSF (20%) decreases the proportion of P-P division. The data is presented as mean fold change  $\pm$  SEM relative to Control IgG (anti-GATA6) (Control IgG = 1.0; NAb =  $0.62 \pm 0.19$ ; LIFR block =  $0.56 \pm 0.13$ ; Control versus LIF NAb, p = n.s.; Control versus LIFR block, p < 0.05; n = 3 with approximately 100 pairs counted per condition; and Mann-Whitney).

(I) Silver staining shows distinct protein patterns in E10.5 CSF collected from lateral versus fourth ventricles. The arrowheads denote protein bands differentially detected in lateral ventricle CSF (red) versus fourth ventricle CSF (blue).

See also Figure S2.

The LIF signaling pathway represents one compelling example of signaling activity in AF and nascent CSF. LIF regulates SOX2 expression in stem cells (Foshay and Gallicano, 2008; Niwa et al., 2009; Onishi and Zandstra, 2015; Pitman et al., 2004; Shimazaki et al., 2001). LIF also stimulates neural progenitor proliferation (Hatta et al., 2002) and influences cell fate at later developmental stages (Onishi and Zandstra, 2015). Thus, we tested if CSF-LIF mediated signaling regulates neural stem cells following neural tube closure. We observed membrane-associated LIFR and phospho-STAT3 in P-Vimentin-positive apical progenitor cells

at E10.5 (Figures 6A and 6B). In agreement with a previous report showing LIF availability in the CSF at E11 (Hatta et al., 2006), we detected LIF-reactive bands in E10.5 CSF treated with the deglycosylating enzyme PNGase F (Figure 6C). Direct injection of recombinant LIF into E10.5 telencephalic ventricles ex vivo stimulated P-STAT3 activity in neuroepithelial progenitor cells (Figure 6D). We then used a pair-cell assay to test for the functional effects of LIF on E10.5 neural progenitor self-renewal (Figure 6E). Using SOX2 to identify neural progenitors and TUJ1 to identify neurons, we found that LIF stimulated self-renewal

(progenitor-progenitor division; P-P) of SOX2-positive neural progenitors at the expense of differentiation (progenitor-neuron division; P-N) or transient cells that co-express SOX2 and TUJ1 (Transient-Transient division; Tr-Tr; Hutton and Pevny, 2011; Figure 6F). Pair-cell assays using 20% E10.5 CSF revealed a similar ability to promote progenitor self-renewal (Figure 6G). The frequency of SOX2-positive P-P divisions declined in cells cultured with CSF containing LIFR blocking or LIF neutralization (NAb) antibodies (Figure 6H), showing that CSF-based signaling via the LIFR can promote neuroepithelial cell self-renewal, though other CSF-borne signals likely work in concert *in vivo* to regulate this process.

*Lifr*-deficient mice are microcephalic by E12.5, a time when LIF is beginning to be produced by the choroid plexus (Gregg and Weiss, 2005). Because the choroid plexus is not yet developed at E10.5, our findings suggest that CSF-LIF must originate from another source to contribute to progenitor proliferation at this earlier developmental stage. While we cannot rule out a contribution by the neuroepithelium (Figure S2E), LIF is abundantly expressed by the amniotic sac and yolk sac (Figure S2E). Consistent with CSF drawing fluid from the AF at this stage (Alonso et al., 1998), 10 kDa fluorescein-conjugated dextrans injected into the intra-amniotic space of E10.5 mouse embryos were detected in the CSF 30 min following injection (average relative fluorescent units: 10 kDa dextran = 445 and 153.3 and Vehicle = 21.3; 33.7; and 50.1). These data suggest that factors available in the AF—possibly including LIF—can reach the CSF by classic diffusion through the neuroepithelium at this age. The zebrafish neuroepithelium is permeable to fluorescent dyes up to 70 kDa (Chang and Sive, 2012).

Our proteomic and functional analyses were performed using pooled samples of AF and CSF. However, silver staining analyses of lateral and fourth ventricle CSF revealed that the CSF proteome was regionalized, with distinct protein bands detected in each sample already at E10.5 (Figure 6I), prior to choroid plexus-driven regionalization that occurs at later ages (Lun et al., 2015a). These data suggest differential diffusion or secretion of proteins into each CSF-filled ventricular subcompartment, and potentially, differential effects on local progenitor progression.

### Quantitative Proteomic Analysis of Maturing AF


We have limited knowledge regarding the mouse AF proteome and how it develops over time. Based upon the observation that E10.5 AF supports the developing olfactory placode (Figure 3E), we performed quantitative MS to reveal changes in the AF proteome as it develops from E8.5 to E10.5 and E14.5. The composition of E10.5 AF and E14.5 AF were less complex than E8.5 AF (Figure 7A; Table S2). Only 80 proteins found in the older AF samples were not detected at E8.5 (Figure 7B; Table S2). Conversely, 476 proteins were exclusive to E8.5 AF (Figure 7B; Table S2). Unsupervised clustering of expressed proteins followed by enrichment analysis revealed that many proteins enriched in E8.5 AF were ribosomal proteins and translation factors (Figures 7C and 7D; Table S2; cluster 1 and 515 members). E14.5 AF was enriched in proteins that were secreted (e.g., collagens) or important for cell adhesion (e.g., NCAM1 and VCAM1) (Figures 7C and 7D; Table S2; cluster 2 and 88 members). Cluster 3 (88 members) hosted proteins that were more abundant at

later ages than at E8.5 (Figure 7C; Table S2). Many of these proteins were secreted (e.g., DKK3 and IGF2) or ECM proteins (e.g., collagens and fibronectin; Figure 7D; Table S2). Only 37 proteins were enriched in E10.5 AF (Figure 7C; Table S2; cluster 5) and some of these (e.g., GAS1, MSTN, and TGFBR3) were members of canonical signaling pathways.

Despite the expansion of the brain's ventricles between E8.5–E10.5 (Figure 1), direct comparison of the AF and CSF proteomes revealed that only 14 proteins found in E10.5 AF were not detected in E10.5 CSF (Figure 7E). These data are consistent with the model that the CSF draws fluid and proteins from the AF (see above; Alonso et al., 1998; Chang and Sive, 2012). Conversely, many proteins were enriched in the E10.5 CSF versus E10.5 AF, including SHH (mean SHH concentration [pg/ml] represented as mean  $\pm$  SEM at E10.5: AF = 50.6  $\pm$  11.9; CSF = 197.6  $\pm$  12.7; n = 3; p < 0.01, and t test; see also Figure 5A). Importantly, 264 proteins in E10.5 CSF were not detected in E10.5 AF, indicating other sources including the adjacent neuroepithelium (see Marzesco et al., 2005) as major contributors to the nascent CSF proteome (Figure 7E). By E14.5, the CSF and AF were even more distinct from each other, with 131 proteins exclusively detected in E14.5 CSF and 110 proteins exclusively detected in E14.5 AF (Figure 7F), reflecting progressive, age-dependent divergence of the AF and CSF. Principal component analysis showed that E8.5 AF and E14.5 CSF were the most distinct fluid compartments, whereas E10.5 CSF, E10.5 AF, and E14.5 AF were more similar to each other (Figure 7G). Thus, the developing AF and CSF undergo a progressive differentiation with instructive capacities matched to adjacent neurogenic tissues.

### DISCUSSION

We present a resource in which we have: (1) developed tools to collect AF and CSF before and after neural tube closure, (2) quantified ventricular expansion in live mouse embryos before and after neural tube closure, (3) systematically documented the dynamic signaling capacity and protein content of the AF and CSF proteomes as they differentiate over the course of embryonic development, (4) developed robust, heterochronic *ex vivo* explant systems to assess functional, instructive activities of AF and nascent CSF in promoting neural progenitor cell fate in the developing neural plate, neural tube, and brain as well as the neurogenic cranial placodes, and (5) demonstrated that CSF-based signaling via the LIFR stimulates neuroepithelial cell self-renewal. Paired with the dynamic morphological features of the early developing head and brain, our findings should accelerate future studies investigating the regulation of fluid/tissue interactions and their consequences for signaling during these early, understudied stages of brain development. Since many tissues of the developing embryo, particularly the precursors of the central and peripheral nervous system, are in contact with AF/CSF, our proteomic data set serves as an important resource for investigators across multiple developmental fields who are interested in studying fluid-tissue interactions and related mechanisms during mid-gestation when embryos are particularly vulnerable to the consequences of disrupted signaling.



**Figure 7. MS Reveals Decreasing Complexity of AF as Embryo Develops from E8.5 to E14.5**

(A) Heatmap of normalized spectral counts reveals differential protein availability in AF between E8.5, E10.5, and E14.5. The unsupervised hierarchical clustering grouped biological replicates of each fluid together (E8.5: n = 3; E10.5: n = 2; and E14.5: n = 2). Each replicate contains 30 µg total AF protein pooled from multiple embryos. The spectral counts were scaled and analyzed as in Figure 4A.

(B) Total of 844 proteins identified across all AF samples are represented as in Figure 4B.

(C) Unsupervised clustering was performed and data are presented as in Figure 4C. There are 38 proteins that showed less than 2-fold difference in availability between the ages and were not included in the clustering.

(D) Functional annotation clustering of proteins in GProX clusters 1, 2, and 3 was performed and is presented as in Figure 4D.

(E) At E10.5, 264 and 14 proteins are exclusive to CSF and AF, respectively, and 240 proteins are common to both fluid compartments.

(legend continued on next page)

## Analyzing Fluid Proteomes and Their Activities in Early Embryos

We addressed several challenges in developing this resource for understanding AF and CSF function in mammalian embryos. A number of studies have used proteomics tools to analyze the AF or CSF at older ages or in other species (see Chang et al., 2015; Dislich et al., 2015; Dziegielewska et al., 1981; Lehtinen et al., 2011; Lun et al., 2015a; Parada et al., 2005; Zappaterra et al., 2007). However, to our knowledge, analysis of the murine AF proteome over the course of early development has not been undertaken, likely due to challenges of reliably collecting small amounts of AF and CSF from early mammalian embryos. We minimized variability in AF collection and preparation, with careful dissection, pooling samples across litters, and assessment of contamination by embryonic or maternal tissues. We similarly standardized CSF collection from embryos after neural tube closure and emergence of distinct brain vesicles. While the CSF between ventricles is regionalized by E10.5 (Figure 6I), analysis of total CSF samples from individual embryos using silver stained gels indicated basic uniformity within and across litters (data not shown). MS analyses showed quantitative differences between pooled AF and CSF samples, which may indicate biological variability independent of technical issues or subtle differences in embryo stages when fluids were collected. Some variability suggested by MS may not reach a threshold for impacting signaling capacity. Quantitative MS may also be limited in its ability to detect low abundance proteins, proteins that have been processed into small peptides, and proteins with other physicochemical properties (e.g., hydrophobicity). However, our analyses of cardinal signaling activity (e.g., BMPs, RA, and SHH) support the model that overall AF and CSF signaling activity is stable among pooled samples. We also detected some variation in AF and CSF activity on neural progenitors in our explant assays; however, these differences may reflect details of the reporter allele (Arnold et al., 2011; Ellis et al., 2004) rather than distinctions of AF or CSF activity on the neuroepithelium. Pair-cell analyses of LIF, a likely AF/CSF signal, indicate stage-specific consistency. Future studies analyzing the influence of AF/CSF on neural stem cell identities in functional assays will further elucidate the biological significance of variability in AF and CSF proteomes.

## The Signaling Capacity of the AF and CSF

The AF and nascent CSF have dynamic morphogenetic signaling capacity via BMPs, RA, and SHH. The effective activity of these signals varies in register with patterning in the embryo, including the brain and the neurogenic cranial placodes. The decrease in total BMP activity during the transition from E8.5 AF to E10.5 CSF may indicate changes in inhibition. For example, FSTL1, a BMP4 antagonist (Geng et al., 2011), was more abundant in E10.5 CSF than in E8.5 AF, while BMP4 was identified in each fluid compartment. The timed delivery of fluid-based signals is also demonstrated by the delayed availability of RA in the

E14.5 CSF, presumably protecting the developing brain from posteriorizing cues, while permitting local signaling via cranial mesenchymal sources once the neural tube closes (Moody and LaMantia, 2015). Finally, our findings of elevated SHH in E10.5 CSF suggest that the developing dorsal forebrain, as well as the ventral forebrain that is thought to receive SHH signals from local tissue sources (Chiang et al., 1996; Rallu et al., 2002b), is exposed to SHH quite early in embryogenesis. In all cases, the action of CSF-borne signals on target cells relies on appropriate receptor expression and ability to transduce signals delivered by the CSF.

Since no individual tissue is solely responsible for the production of AF and nascent CSF at the early ages examined (e.g., E8.5 and E10.5; the choroid plexus contributes to CSF production in our E14.5 samples), there is no transcriptome from which to derive the early AF and CSF proteomes. Therefore, our MS approach provides a singular directory of hundreds of differentially available proteins in the maturing AF and CSF over time. The diversity of proteins identified in early AF and CSF includes growth factors and their receptors (e.g., BOC and GAS1 for SHH), axon guidance molecules (e.g., semaphorins, DCC, and ephrins), and ECM components (e.g., reelin and laminin subunits). The presence of prominin-1-containing membrane particles (Dubreuil et al., 2007; Marzesco et al., 2005) in E10.5 and E14.5 CSF, but not E8.5 AF (Table S1), suggest that the developing neuroepithelium contributes to the CSF proteome. Whether these particles have an active role in CSF-based signaling (Cossetti et al., 2014; Feliciano et al., 2014) or are cellular waste products cleared from the developing neural tissue, remains to be determined. Nevertheless, signaling components of many of the pathways represented in the AF and CSF proteomes have roles in neural tube closure (Wilde et al., 2014), inviting hypotheses for fluid-based regulation of this developmental stage.

## AF and CSF Proteomes and Hypotheses for Early Mammalian Brain Development

The resources of AF/CSF proteomes, activities, and embryo-based assays to evaluate signaling capacities also provide an essential foundation for generating and testing hypotheses of mammalian neural development and how it is compromised by genetic and environmental disruption. We found that AF and CSF vary in several dimensions within 48 hr during early to mid-stage mouse embryogenesis. Changes in osmolality and fluid volumes may have substantial consequences for morphogenetic movements that maintain the neural plate and then support neural tube closure. Importantly, distinctions in AF/CSF protein constituents—especially metabolic and receptor proteins in the fluids at different stages—suggest key changes in cells of embryonic surface tissues that directly contact AF or CSF during the transition from neural plate to forebrain vesicle epithelium. Such changes may be driven by AF versus CSF proteomes and may be metabolic indicators of molecular and

(F) By E14.5, 131 and 110 proteins are exclusive to CSF and AF, respectively, and 207 proteins are shared between the two fluid compartments.

(G) Principal component analysis of protein spectral counts revealed greater degree of variance between fluid compartments than between biological replicates within each compartment. The E8.5 AF and E14.5 CSF were the most distinct fluid compartments, whereas E10.5 CSF, E10.5 AF, and E14.5 AF were more similar to each other. PC1 and PC2 explained 48.9% and 24.5% of the total variance, respectively. The ellipses and lines indicate the 95% confidence interval. See also Table S2.

cellular distinctions in embryonic tissues that influence stem cell progression and early neurogenesis. Despite tremendous focus on neural stem cell progression at later stages of brain development in the spinal cord, hindbrain, and forebrain (see [Dias et al., 2014](#)), there has been far less attention paid to the key transition between early neuroectodermal stem cell and neural stem cell identity as the neural tube closes. Mining of our proteomic resource, combined with use of our assays to evaluate stem cell progression will provide important insights into mechanisms that mediate this key period of mammalian brain development.

The early AF and CSF proteomes reflect, in varying degrees, the balance between maternally derived protein constituents and other small molecules and those derived from the embryo. These proteomes are likely altered dramatically in response to maternal illness, changes in nutrition, substance abuse, and exposure to environmental toxins ([Wilde et al., 2014](#)). Thus, the definition of dynamic AF and CSF activities we have provided constitutes an essential resource for understanding risk and consequence of maternal/fetal interactions during first trimester development.

## EXPERIMENTAL PROCEDURES

### Animals

Timed pregnant CD1 dams were obtained from Charles River Laboratories. *Sox2:EGFP* ([Ellis et al., 2004](#)) was shared by L. Pevny. All animal experimentation was carried out under protocols approved by the Institutional Animal Care and Use Committees of Boston Children's Hospital and The George Washington University School of Medicine.

### Ventricular System Measurements

Uterine horns were placed on Sylgard plates, immersed in PBS, and ultrasound image sequences were captured (Vivo 2100, Visual Sonics; MS550 transducer; step size: 0.076 mm). 3D reconstruction of ventricular system was performed for visualization (Vivo 2100) and quantification (Fiji-TrakEM).

### AF and CSF Collection and Measurements

AF was collected by inserting a glass capillary into the intra-amniotic space. CSF was collected by inserting a glass capillary into telencephalic ventricles, aqueduct, fourth ventricle, or cisterna magna, and processed as described ([Zappaterra et al., 2013](#)). Protein concentration (BCA kit) and osmolality (10 µl fluid input, Vapro 5600) were measured.

### Forebrain Neuroectodermal Explants

E8.5 anterior neuroectoderm containing the presumptive forebrain was dissected from the anterior neural ridge to the optic vesicle. E10.5 dorsal telencephalic neuroepithelium was isolated by: (1) dissecting the cortical neuroepithelium caudal to the olfactory bulb at the anterior boundary of the lateral ganglionic eminence (LGE), (2) dissecting caudal to the posterior boundary of the LGE and extending to the medial wall of the developing cortical rudiment, (3) separating the LGE from the developing cortex, and (4) removing the midline at the apex of the neocortex where the lateral cortical surface meets the interhemispheric wall. Following 30 min of enzymatic digestion with pancreatin on ice, neuroepithelium was separated from mesenchyme and surface ectoderm using tungsten needles. Explants were placed on polycarbonate membranes (8.0 µm pore size; Whatman) in imaging dishes (MatTek), and cultured in 20% AF or CSF in Neurobasal medium supplemented with antibiotics and L-glutamine. Explants were cultured for 24 hr.

## SUPPLEMENTAL INFORMATION

Supplemental Information includes Supplemental Experimental Procedures, two figures, and two tables and can be found with this article online at <http://dx.doi.org/10.1016/j.devcel.2015.11.015>.

## AUTHOR CONTRIBUTIONS

K.F.C., M.W.S., H.S., A.S.L., and M.K.L. designed research; K.G.B. and H.S. developed proteomic tools and analysis pipeline; K.F.C., M.W.S., K.G.B., H.P., S.T., M.P.L., H.M., T.M., H.S., A.S.L., and M.K.L. performed research; K.F.C., M.W.S., K.G.B., H.P., S.T., T.M., H.S., A.S.L., and M.K.L. analyzed data; K.F.C. and M.K.L. wrote the manuscript; and all authors edited the manuscript.

## ACKNOWLEDGMENTS

We thank C. Walsh for helpful discussions; E. Paronett, T. Adelita, and A. Malesz for technical assistance; and J. Steen for RPS12 and RPL11 antibodies. We are grateful for the following support: National Science Foundation Graduate Research Fellowship (K.F.C.); NIH R01 DC011534 (A.S.L.); and Pediatric Hydrocephalus Foundation, Eleanor and Miles Shore Fellowship Program for Scholars in Medicine/Boston Children's Hospital Career Development Award, Alfred P. Sloan Foundation, BCH IDDRC P30 HD18655, NIH K99/R00 NS072192, and R01 NS088566 (M.K.L.).

Received: May 12, 2015

Revised: September 28, 2015

Accepted: November 16, 2015

Published: December 21, 2015

## REFERENCES

- [Alonso, M.I., Gato, A., Moro, J.A., and Barbosa, E. \(1998\). Disruption of proteoglycans in neural tube fluid by beta-D-xyloside alters brain enlargement in chick embryos. \*Anat. Rec.\* 252, 499–508.](#)
- [Arbeille, E., Reynaud, F., Sanyas, I., Bozon, M., Kindbeiter, K., Causeret, F., Pierani, A., Falk, J., Moret, F., and Castellani, V. \(2015\). Cerebrospinal fluid-derived Semaphorin3B orients neuroepithelial cell divisions in the apicobasal axis. \*Nat. Commun.\* 6, 6366.](#)
- [Arnold, K., Sarkar, A., Yram, M.A., Polo, J.M., Bronson, R., Sengupta, S., Seandel, M., Geijsen, N., and Hochedlinger, K. \(2011\). Sox2\(+\) adult stem and progenitor cells are important for tissue regeneration and survival of mice. \*Cell Stem Cell\* 9, 317–329.](#)
- [Bjornsson, C.S., Apostolopoulou, M., Tian, Y., and Temple, S. \(2015\). It takes a village: constructing the neurogenic niche. \*Dev. Cell\* 32, 435–446.](#)
- [Chang, J.T., and Sive, H. \(2012\). An assay for permeability of the zebrafish embryonic neuroepithelium. \*J. Vis. Exp.\* 42, e4242.](#)
- [Chang, J.T., Lehtinen, M.K., and Sive, H. \(2015\). Zebrafish cerebrospinal fluid mediates cell survival through a retinoid signaling pathway. \*Dev. Neurobiol.\* <http://dx.doi.org/10.1002/dneu.22300>.](#)
- [Chatzci, C., Cunningham, T.J., and Duester, G. \(2013\). Investigation of retinoic acid function during embryonic brain development using retinaldehyde-rescued Rdh10 knockout mice. \*Dev. Dyn.\* 242, 1056–1065.](#)
- [Chiang, C., Litingtung, Y., Lee, E., Young, K.E., Corden, J.L., Westphal, H., and Beachy, P.A. \(1996\). Cyclopia and defective axial patterning in mice lacking Sonic hedgehog gene function. \*Nature\* 383, 407–413.](#)
- [Cossetti, C., Iraci, N., Mercer, T.R., Leonardi, T., Alpi, E., Drago, D., Alfaro-Cervello, C., Saini, H.K., Davis, M.P., Schaeffer, J., et al. \(2014\). Extracellular vesicles from neural stem cells transfer IFN- \$\gamma\$  via Ifngr1 to activate Stat1 signaling in target cells. \*Mol. Cell\* 56, 193–204.](#)
- [Desmond, M.E., and Jacobson, A.G. \(1977\). Embryonic brain enlargement requires cerebrospinal fluid pressure. \*Dev. Biol.\* 57, 188–198.](#)
- [Dias, J.M., Alekseenko, Z., Applequist, J.M., and Ericson, J. \(2014\). Tgf \$\beta\$  signaling regulates temporal neurogenesis and potency of neural stem cells in the CNS. \*Neuron\* 84, 927–939.](#)
- [Dislich, B., Wohlrab, F., Bachhuber, T., Mueller, S., Kuhn, P.H., Hogl, S., Meyer-Luehmann, M., and Lichtenthaler, S.F. \(2015\). Label-free quantitative proteomics of mouse cerebrospinal fluid detects BACE1 protease substrates in vivo. \*Mol. Cell. Proteomics\* 14, 2550–2563.](#)

- Doan, L.T., Javier, A.L., Furr, N.M., Nguyen, K.L., Cho, K.W., and Monuki, E.S. (2012). A Bmp reporter with ultrasensitive characteristics reveals that high Bmp signaling is not required for cortical hem fate. *PLoS ONE* 7, e44009.
- Dubreuil, V., Marzesco, A.M., Corbeil, D., Huttner, W.B., and Wilsch-Bräuninger, M. (2007). Midbody and primary cilium of neural progenitors release extracellular membrane particles enriched in the stem cell marker prominin-1. *J. Cell Biol.* 176, 483–495.
- Durston, A.J., Timmermans, J.P., Hage, W.J., Hendriks, H.F., de Vries, N.J., Heideveld, M., and Nieuwkoop, P.D. (1989). Retinoic acid causes an antero-posterior transformation in the developing central nervous system. *Nature* 340, 140–144.
- Dziegielewska, K.M., Evans, C.A., Lai, P.C., Lorscheider, F.L., Malinowska, D.H., Möllgård, K., and Saunders, N.R. (1981). Proteins in cerebrospinal fluid and plasma of fetal rats during development. *Dev. Biol.* 83, 193–200.
- Ek, C.J., Dziegielewska, K.M., and Saunders, N.R. (2005). Development of the blood-cerebrospinal fluid barrier. In *The Blood-Cerebrospinal Fluid Barrier*, W. Zheng and A. Chodobski, eds. (Boca Raton: Taylor & Francis), pp. 3–23.
- Ellis, P., Fagan, B.M., Magness, S.T., Hutton, S., Taranova, O., Hayashi, S., McMahon, A., Rao, M., and Pevny, L. (2004). SOX2, a persistent marker for multipotential neural stem cells derived from embryonic stem cells, the embryo or the adult. *Dev. Neurosci.* 26, 148–165.
- Feliciano, D.M., Zhang, S., Nasrallah, C.M., Lisgo, S.N., and Bordey, A. (2014). Embryonic cerebrospinal fluid nanovesicles carry evolutionarily conserved molecules and promote neural stem cell amplification. *PLoS ONE* 9, e88810.
- Foshay, K.M., and Gallicano, G.I. (2008). Regulation of Sox2 by STAT3 initiates commitment to the neural precursor cell fate. *Stem Cells Dev.* 17, 269–278.
- Gato, A., and Desmond, M.E. (2009). Why the embryo still matters: CSF and the neuroepithelium as interdependent regulators of embryonic brain growth, morphogenesis and histogenesis. *Dev. Biol.* 327, 263–272.
- Geng, Y., Dong, Y., Yu, M., Zhang, L., Yan, X., Sun, J., Qiao, L., Geng, H., Nakajima, M., Furuichi, T., et al. (2011). Follistatin-like 1 (Fstl1) is a bone morphogenetic protein (BMP) 4 signaling antagonist in controlling mouse lung development. *Proc. Natl. Acad. Sci. USA* 108, 7058–7063.
- Gregg, C., and Weiss, S. (2005). CNTF/LIF/gp130 receptor complex signaling maintains a VZ precursor differentiation gradient in the developing ventral forebrain. *Development* 132, 565–578.
- Grove, E.A., and Monuki, E.S. (2013). Morphogens, patterning centers, and their mechanisms of action. In *Patterning and Cell Type Specification in the Developing CNS and PNS*, J.L. Rubenstein and P. Rakic, eds. (San Diego: Elsevier), pp. 25–44.
- Haskell, G.T., and LaMantia, A.S. (2005). Retinoic acid signaling identifies a distinct precursor population in the developing and adult forebrain. *J. Neurosci.* 25, 7636–7647.
- Hatta, T., Moriyama, K., Nakashima, K., Taga, T., and Otani, H. (2002). The Role of gp130 in cerebral cortical development: in vivo functional analysis in a mouse exo utero system. *J. Neurosci.* 22, 5516–5524.
- Hatta, T., Matsumoto, A., Ono, A., Udagawa, J., Nimura, M., Hashimoto, R., and Otani, H. (2006). Quantitative analyses of leukemia inhibitory factor in the cerebrospinal fluid in mouse embryos. *Neuroreport* 17, 1863–1866.
- Higginbotham, H., Guo, J., Yokota, Y., Umberger, N.L., Su, C.Y., Li, J., Verma, N., Hirt, J., Ghukasyan, V., Caspary, T., and Anton, E.S. (2013). Arl13b-regulated cilia activities are essential for polarized radial glial scaffold formation. *Nat. Neurosci.* 16, 1000–1007.
- Huang, W., Sherman, B.T., and Lempicki, R.A. (2009a). Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat. Protoc.* 4, 44–57.
- Huang, W., Sherman, B.T., and Lempicki, R.A. (2009b). Bioinformatics enrichment tools: paths toward the comprehensive functional analysis of large gene lists. *Nucleic Acids Res.* 37, 1–13.
- Huang, X., Ketova, T., Fleming, J.T., Wang, H., Dey, S.K., Litingtung, Y., and Chiang, C. (2009c). Sonic hedgehog signaling regulates a novel epithelial progenitor domain of the hindbrain choroid plexus. *Development* 136, 2535–2543.
- Huang, X., Liu, J., Ketova, T., Fleming, J.T., Grover, V.K., Cooper, M.K., Litingtung, Y., and Chiang, C. (2010). Transventricular delivery of Sonic hedgehog is essential to cerebellar ventricular zone development. *Proc. Natl. Acad. Sci. USA* 107, 8422–8427.
- Hunter, K.E., and Hatten, M.E. (1995). Radial glial cell transformation to astrocytes is bidirectional: regulation by a diffusible factor in embryonic forebrain. *Proc. Natl. Acad. Sci. USA* 92, 2061–2065.
- Hutton, S.R., and Pevny, L.H. (2011). SOX2 expression levels distinguish between neural progenitor populations of the developing dorsal telencephalon. *Dev. Biol.* 352, 40–47.
- Kokovay, E., Goderie, S., Wang, Y., Lotz, S., Lin, G., Sun, Y., Roysam, B., Shen, Q., and Temple, S. (2010). Adult SVZ lineage cells home to and leave the vascular niche via differential responses to SDF1/CXCR4 signaling. *Cell Stem Cell* 7, 163–173.
- LaMantia, A.S., Colbert, M.C., and Linney, E. (1993). Retinoic acid induction and regional differentiation prefigure olfactory pathway formation in the mammalian forebrain. *Neuron* 10, 1035–1048.
- LaMantia, A.S., Bhasin, N., Rhodes, K., and Heemskerk, J. (2000). Mesenchymal/epithelial induction mediates olfactory pathway formation. *Neuron* 28, 411–425.
- Layton, M.J., Cross, B.A., Metcalf, D., Ward, L.D., Simpson, R.J., and Nicola, N.A. (1992). A major binding protein for leukemia inhibitory factor in normal mouse serum: identification as a soluble form of the cellular receptor. *Proc. Natl. Acad. Sci. USA* 89, 8616–8620.
- Lehtinen, M.K., and Walsh, C.A. (2011). Neurogenesis at the brain-cerebrospinal fluid interface. *Annu. Rev. Cell Dev. Biol.* 27, 653–679.
- Lehtinen, M.K., Zappaterra, M.W., Chen, X., Yang, Y.J., Hill, A.D., Lun, M., Maynard, T., Gonzalez, D., Kim, S., Ye, P., et al. (2011). The cerebrospinal fluid provides a proliferative niche for neural progenitor cells. *Neuron* 69, 893–905.
- Lowery, L.A., and Sive, H. (2009). Totally tubular: the mystery behind function and origin of the brain ventricular system. *BioEssays* 31, 446–458.
- Lun, M.P., Johnson, M.B., Broadbelt, K.G., Watanabe, M., Kang, Y.-J., Chau, K.F., Springel, M.W., Malesz, A., Sousa, A.M.M., Pletikos, M., et al. (2015a). Spatially heterogeneous choroid plexus transcriptomes encode positional identity and contribute to regional CSF production. *J. Neurosci.* 35, 4903–4916.
- Lun, M.P., Monuki, E.S., and Lehtinen, M.K. (2015b). Development and functions of the choroid plexus-cerebrospinal fluid system. *Nat. Rev. Neurosci.* 16, 445–457.
- Manzini, M.C., and Walsh, C.A. (2011). What disorders of cortical development tell us about the cortex: one plus one does not always make two. *Curr. Opin. Genet. Dev.* 21, 333–339.
- Marzesco, A.M., Janich, P., Wilsch-Bräuninger, M., Dubreuil, V., Langenfeld, K., Corbeil, D., and Huttner, W.B. (2005). Release of extracellular membrane particles carrying the stem cell marker prominin-1 (CD133) from neural progenitors and other epithelial cells. *J. Cell Sci.* 118, 2849–2858.
- Massarwa, R., and Niswander, L. (2013). *In toto* live imaging of mouse morphogenesis and new insights into neural tube closure. *Development* 140, 226–236.
- McConnell, S.K., and Kaznowski, C.E. (1991). Cell cycle dependence of laminar determination in developing neocortex. *Science* 254, 282–285.
- Moody, S.A., and LaMantia, A.S. (2015). Transcriptional regulation of cranial sensory placode development. *Curr. Top. Dev. Biol.* 111, 301–350.
- Niwa, H., Ogawa, K., Shimosato, D., and Adachi, K. (2009). A parallel circuit of LIF signalling pathways maintains pluripotency of mouse ES cells. *Nature* 460, 118–122.
- Onishi, K., and Zandstra, P.W. (2015). LIF signaling in stem cells and development. *Development* 142, 2230–2236.
- Parada, C., Gato, A., and Bueno, D. (2005). Mammalian embryonic cerebrospinal fluid proteome has greater apolipoprotein and enzyme pattern complexity than the avian proteome. *J. Proteome Res.* 4, 2420–2428.
- Pitman, M., Emery, B., Binder, M., Wang, S., Butzkueven, H., and Kilpatrick, T.J. (2004). LIF receptor signaling modulates neural stem cell renewal. *Mol. Cell. Neurosci.* 27, 255–266.

- Rallu, M., Corbin, J.G., and Fishell, G. (2002a). Parsing the prosencephalon. *Nat. Rev. Neurosci.* 3, 943–951.
- Rallu, M., Machold, R., Gaiano, N., Corbin, J.G., McMahon, A.P., and Fishell, G. (2002b). Dorsoventral patterning is established in the telencephalon of mutants lacking both Gli3 and Hedgehog signaling. *Development* 129, 4963–4974.
- Rawson, N.E., Lischka, F.W., Yee, K.K., Peters, A.Z., Tucker, E.S., Meehan, D.W., Zirlinger, M., Maynard, T.M., Burd, G.B., Dulac, C., et al. (2010). Specific mesenchymal/epithelial induction of olfactory receptor, vomeronasal, and gonadotropin-releasing hormone (GnRH) neurons. *Dev. Dyn.* 239, 1723–1738.
- Rigbolt, K.T., Vanselow, J.T., and Blagoev, B. (2011). GProX, a user-friendly platform for bioinformatics analysis and visualization of quantitative proteomics data. *Mol. Cell Proteomics* 10, <http://dx.doi.org/10.1074/mcp.O110.007450>.
- Sawamoto, K., Wichterle, H., Gonzalez-Perez, O., Cholfin, J.A., Yamada, M., Spassky, N., Murcia, N.S., Garcia-Verdugo, J.M., Marin, O., Rubenstein, J.L., et al. (2006). New neurons follow the flow of cerebrospinal fluid in the adult brain. *Science* 311, 629–632.
- Shen, Q., Goderie, S.K., Jin, L., Karanth, N., Sun, Y., Abramova, N., Vincent, P., Pumiglia, K., and Temple, S. (2004). Endothelial cells stimulate self-renewal and expand neurogenesis of neural stem cells. *Science* 304, 1338–1340.
- Shen, Q., Wang, Y., Dimos, J.T., Fasano, C.A., Phoenix, T.N., Lemischka, I.R., Ivanova, N.B., Stifani, S., Morrisey, E.E., and Temple, S. (2006). The timing of cortical neurogenesis is encoded within lineages of individual progenitor cells. *Nat. Neurosci.* 9, 743–751.
- Shimazaki, T., Shingo, T., and Weiss, S. (2001). The ciliary neurotrophic factor/leukemia inhibitory factor/gp130 receptor complex operates in the maintenance of mammalian forebrain neural stem cells. *J. Neurosci.* 21, 7642–7653.
- Siegenthaler, J.A., Ashique, A.M., Zarbalis, K., Patterson, K.P., Hecht, J.H., Kane, M.A., Folias, A.E., Choe, Y., May, S.R., Kume, T., et al. (2009). Retinoic acid from the meninges regulates cortical neuron generation. *Cell* 139, 597–609.
- Stavridis, M.P., Collins, B.J., and Storey, K.G. (2010). Retinoic acid orchestrates fibroblast growth factor signalling to drive embryonic stem cell differentiation. *Development* 137, 881–890.
- Tavazoie, M., Van der Veken, L., Silva-Vargas, V., Louissaint, M., Colonna, L., Zaidi, B., Garcia-Verdugo, J.M., and Doetsch, F. (2008). A specialized vascular niche for adult neural stem cells. *Cell Stem Cell* 3, 279–288.
- Tomida, M. (2000). Structural and functional studies on the leukemia inhibitory factor receptor (LIF-R): gene and soluble form of LIF-R, and cytoplasmic domain of LIF-R required for differentiation and growth arrest of myeloid leukemic cells. *Leuk. Lymphoma* 37, 517–525.
- Tong, C.K., Han, Y.G., Shah, J.K., Obernier, K., Guinto, C.D., and Alvarez-Buylla, A. (2014). Primary cilia are required in a unique subpopulation of neural progenitors. *Proc. Natl. Acad. Sci. USA* 111, 12438–12443.
- Tucker, E.S., Segall, S., Gopalakrishna, D., Wu, Y., Vernon, M., Polleux, F., and Lamantia, A.S. (2008). Molecular specification and patterning of progenitor cells in the lateral and medial ganglionic eminences. *J. Neurosci.* 28, 9504–9518.
- Tucker, E.S., Lehtinen, M.K., Maynard, T., Zirlinger, M., Dulac, C., Rawson, N., Pavny, L., and Lamantia, A.-S.S. (2010). Proliferative and transcriptional identity of distinct classes of neural precursors in the mammalian olfactory epithelium. *Development* 137, 2471–2481.
- Vermot, J., Gallego Llamas, J., Fraulob, V., Niederreither, K., Chambon, P., and Dollé, P. (2005). Retinoic acid controls the bilateral symmetry of somite formation in the mouse embryo. *Science* 308, 563–566.
- Warde-Farley, D., Donaldson, S.L., Comes, O., Zuberi, K., Badrawi, R., Chao, P., Franz, M., Grouios, C., Kazi, F., Lopes, C.T., et al. (2010). The GeneMANIA prediction server: biological network integration for gene prioritization and predicting gene function. *Nucleic Acids Res.* 38, W214–W220.
- Wilde, J.J., Petersen, J.R., and Niswander, L. (2014). Genetic, epigenetic, and environmental contributions to neural tube closure. *Annu. Rev. Genet.* 48, 583–611.
- Yamamoto, M., McCaffery, P., and Dräger, U.C. (1996). Influence of the choroid plexus on cerebellar development: analysis of retinoic acid synthesis. *Brain Res. Dev. Brain Res.* 93, 182–190.
- Yeh, C., Li, A., Chuang, J.Z., Saito, M., Cáceres, A., and Sung, C.H. (2013). IGF-1 activates a cilium-localized noncanonical G $\beta$  signaling pathway that regulates cell-cycle progression. *Dev. Cell* 26, 358–368.
- Zappaterra, M.W., and Lehtinen, M.K. (2012). The cerebrospinal fluid: regulator of neurogenesis, behavior, and beyond. *Cell. Mol. Life Sci.* 69, 2863–2878.
- Zappaterra, M.D., Lisgo, S.N., Lindsay, S., Gygi, S.P., Walsh, C.A., and Ballif, B.A. (2007). A comparative proteomic analysis of human and rat embryonic cerebrospinal fluid. *J. Proteome Res.* 6, 3537–3548.
- Zappaterra, M.W., LaMantia, A.S., Walsh, C.A., and Lehtinen, M.K. (2013). Isolation of cerebrospinal fluid from rodent embryos for use with dissected cerebral cortical explants. *J. Vis. Exp.* 11, e50333.